ECONSTOR
Make Your Publication Visible

A Service of

ZBW Leibniz-Informationszentrum Wirtschaft
Leibniz Information Centre for Economics

Nie, Peng; Sousa-Poza, Alfonso; Xue, Jianhong

Working Paper

Fuel for life: Domestic cooking fuels and women's health in rural China?

Hohenheim Discussion Papers in Business, Economics and Social Sciences, No. 08-2016

Provided in Cooperation with:
Faculty of Business, Economics and Social Sciences, University of Hohenheim

Suggested Citation: Nie, Peng; Sousa-Poza, Alfonso; Xue, Jianhong (2016) : Fuel for life: Domestic cooking fuels and women's health in rural China?, Hohenheim Discussion Papers in Business, Economics and Social Sciences, No. 08-2016, http://nbn-resolving.de/urn:nbn:de:bsz:100-opus-12356

This Version is available at:
http://hdl.handle.net/10419/142691

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

WWW.ECONSTOR.EU

Mitglied der Leibniz-Gemeinschaft

UNIVERSITY OF HOHENHEIM
FACULTY OF BUSINESS, ECONOMICS AND SOCIAL SCIENCES

HOHENHEIM DISCUSSION PAPERS
IN BUSINESS, ECONOMICS AND SOCIAL SCIENCES

Institute of Health Care & Public Management

DISCUSSION PAPER 08-2016

FUEL FOR LIFE: DOMESTIC COOKING FUELS AND WOMEN'S HEALTH IN RURAL CHINA

Peng Nie
University of Hohenheim

Alfonso Sousa-Poza
University of Hohenheim

Jianhong Xue
Northwest A&F University



State: June 2016

www.wiso.uni-hohenheim.de

Discussion Paper 08-2016

Fuel for Life: Domestic Cooking Fuels and Women's Health in Rural China?

Peng Nie, Alfonso Sousa-Poza, Jianhong Xue

Download this Discussion Paper from our homepage:

https://wiso.uni-hohenheim.de/papers

ISSN 2364-2076 (Printausgabe)
ISSN 2364-2084 (Internetausgabe)

Die Hohenheim Discussion Papers in Business, Economics and Social Sciences dienen der schnellen Verbreitung von Forschungsarbeiten der Fakultät Wirtschafts- und Sozialwissenschaften. Die Beiträge liegen in alleiniger Verantwortung der Autoren und stellen nicht notwendigerweise die Meinung der Fakultät Wirtschafts- und Sozialwissenschaften dar.

Hohenheim Discussion Papers in Business, Economics and Social Sciences are intended to make results of the Faculty of Business, Economics and Social Sciences research available to the public in order to encourage scientific discussion and suggestions for revisions. The authors are solely responsible for the contents which do not necessarily represent the opinion of the Faculty of Business, Economics and Social Sciences.

Fuel for Life: Domestic Cooking Fuels and Women's Health in Rural China

PENG NIE[1,*], ALFONSO SOUSA-POZA[1] AND JIANHONG XUE[2]

[1]*Institute for Health Care & Public Management, University of Hohenheim, 70599 Stuttgart, Germany*

[2]*Department of Agricultural Economics, College of Economics and Management, Northwest A&F University, 3 Taicheng Road, Yangling, Shaanxi 712100, China*

ABSTRACT

Using longitudinal and biomarker data from the China Family Panel Studies and the China Health and Nutrition Survey, this study examines the association between the type of domestic cooking fuel and the health of women aged ≥16 in rural China. Regarding three major domestic cooking fuels (wood/straw, coal and liquefied natural gas (LNG)), we find that, compared to women whose households cook with dirty fuels like wood/straw, women whose households cook with cleaner fuels like LNG have a significantly lower probability of chronic or acute diseases and are more likely to report better health. Even after controlling for unobserved individual heterogeneity, we find some evidence that women in households cooking with LNG are less likely to suffer from chronic/acute diseases. Cooking with domestic coal instead of wood or straw is also associated with elevated levels of having certain risks (such as systolic and diastolic blood pressure) related to cardiovascular diseases.

JEL Classification Codes: I10, D10, J10; Q53

Keywords: household cooking fuels; health; women; rural China

* Correspondence: Peng Nie, Email: Peng_Nie@uni-hohenheim.de

Fuel for Life: Domestic Cooking Fuels and Women's Health in Rural China

1. Introduction

Globally, 40% of the population relies on solid fuels, including coal and biomass (e.g. wood, charcoal, agricultural residues and dung), for domestic cooking, thereby breathing in a large amount of particulate matter (PM) and pollutants that can be detrimental to health (United Nations Development Group, 2013). According to the 2010 Global Burden of Disease/Comparative Risk Assessment Project, for example, exposure to indoor air pollution (IAP) from cooking with solid fuels resulted in 3.5 million premature death and various other health problems (e.g. lung cancer) in 2010 (Lim et al., 2012; Reid et al., 2012). Yet despite unprecedented economic development in China, most rural Chinese still cook with solid fuels, 47.6% with biomass and 13.5% with coal in 2011 (Duan et al., 2014), and the use of solid fuels is likely to continue in the future (Zhang and Wu, 2012). As a result, IAP associated with solid fuels is estimated to be the largest single environmental risk contributor and ranks sixth among all risk factors examined for bad health in China (Zhang and Smith, 2007), resulting in over half a million premature deaths annually (World Health Organization and United Nations Development Programme, 2009). At the same time, although a relatively large body of epidemiological and environmental literature examines the effects of IAP on health in parts of China (Finkelman et al., 1999; Smith et al., 2004; Peabody et al., 2005; Zhao et al., 2006; Liu et al., 2007a; Zhang and Smith, 2007; Baumgartner et al., 2011; Kan et al., 2011; Jin et al., 2014), no studies assess this risk using nationally representative data and a longitudinal setting. Consequently, the evidence on IAP and health in China as a whole is scant, often narrowly focused (i.e. on certain provinces or cities) and usually based on cross-sectional analyses with limited generalizability and little ability to address confounders.

To address these shortcomings, this paper investigates the impacts of domestic cooking fuels on the health outcomes of rural women aged ≥16 using 2010–2012 data from the China Family Panel Studies (CFPS) and 1991–2009 data from the China Health and Nutrition Survey (CHNS). Specifically, it analyses the different health impacts associated with three major household cooking fuels: wood/straw, coal and liquefied natural gas (LNG). The research

focuses on women in rural China because they are predominantly responsible for cooking and thus bear the brunt of the illness burdens associated with IAP (Zhang and Smith, 2007).

The study contributes to the existing literature in three ways: First, it provides a comprehensive analysis of the health impacts related to domestic cooking fuels by using the most recent nationally and regionally representative datasets (the CPFS and CHNS, respectively). Second, it assesses the impact of household cooking fuels on health within a panel setting capable of capturing time-invariant individual heterogeneity. Third, by using biomarker data from the 2009 CHNS, it more closely examines the association between IAP and certain risk predictors of cardiovascular disease, evidence for which has been primarily limited by a lack of reliable biomarker data (Fullerton et al., 2008).[1] This association can shed new light on the mechanisms by which IAP from household cooking fuels may influence health outcomes.

The remainder of the discussion proceeds as follows: Section 2 reviews the relevant research on this topic in China. Section 3 outlines the data and methodologies, and Section 4 reports the empirical results. Section 5 concludes the paper.

2. Prior studies

2.1. Fuel-based IAP and respiratory diseases

In the rapidly growing body of epidemiologic and environmental science literature on the association between fuel-based IAP and health in China, most studies present strong evidence that exposure to IAP is significantly related to an increased risk of bad health. With regard to the impact of fuel-based IAP on adult respiratory diseases, Liu et al. (2007b), using survey data for rural Yunyan in Guangdong province, show that cooking with biomass fuel, as compared with liquefied petroleum gas, increases the probability of chronic obstructive pulmonary disease (COPD) in non-smoking women aged ≥40. Similarly, using cross-sectional data for rural China (Shaanxi, Hubei and Zhejiang), Peabody et al. (2005) find that, compared with other types of fuel,[2] coal fuel is associated with an increased level of exhaled CO, an increased history of overall respiratory disease (including asthma, chronic bronchitis, emphysema,

[1] As emphasized by Balakrishnan et al. (2014) and Zhang et al. (2013), biomarkers are considered a very reliable source of information about absorbed doses of pollutants.

[2] Cooking fuels include wood, crop fuel (crop residues and other biomass like dung), coal, charcoal and cleaner fuels (electricity, liquefied petroleum gas and biogas) (Peabody et al., 2005).

COPD and tuberculosis) and decreased forced vital capacity (FVC)[3] among adults aged ≥18. Pan et al. (2002) also show that IAP from fuel combustion for cooking and heating has adverse effects on asthma attacks among respondents aged 15–65. These findings are confirmed by a recent meta-analysis of 33 international studies on IAP's detrimental respiratory impacts among those aged ≥65 (Bentayeb et al., 2013), which points to a generally significant association between IAP exposure and various short-term and long-term respiratory diseases, including wheezing, breathlessness, cough, phlegm, asthma and COPD. In contrast, Xu et al. (2007) find no association between domestic cooking fuels and COPD in individuals aged ≥35 in Nanjing city, while Kan et al.'s (2011) matched case-control analysis of respondents aged ≥15 in Huaiyuan county, Anhui province, identifies no link between cooking with solid fuel and tuberculosis in the presence of proper ventilation. This latter is supported by a recent review of 15 international studies (Lin et al., 2014), which suggests that the association between domestic use of solid fuels and tuberculosis is very low.

Because lung cancer is a serious health problem in China (see She et al., 2013; Chen et al., 2014), many studies focus on its association with fuel-based IAP. For example, one meta-analysis (Zhao et al., 2006) of 27 studies on air pollution and lung cancer risks in China provides evidence that the odds ratio (OR) of the lung cancer risk among women is 1.83 in households that use coal for heating and cooking. Another review by Smith et al. (2004) on the health effect of IAP from coal use in nine single-province/city/municipality studies (seven in China,[4] one in Japan and another in the U.S., with a focus either on women or on men and women separately) finds that the OR of lung cancer risks among women is 1.17. This risk of death from lung cancer for women almost doubles (OR = 1.94) in studies that adjust for smoking behaviour and a history of chronic respiratory disease. A more recent meta-analysis of 23 studies (Kurmi et al., 2012), including 17 carried out in China, also suggests that coal use is correlated with lung cancer and that the risk is more pronounced among females. Kim et al. (2014), using a prospective cohort of women in urban Shanghai from 1996 to 2009, demonstrate that cooking with coal is particularly harmful when the kitchen has poor ventilation, and Barone-Adesi et al. (2012), using a retrospective cohort of 37,272 individuals in Xuanwei county, Yunnan province, show that smoky coal is related to a higher lung cancer risk than smokeless coal. Coal use is also associated with general lung function impairment

[3] FVC is the maximal volume of air exhaled with maximally forced effort from a maximal inspiration; that is, the vital capacity performed with a maximally forced expiratory effort (in liters at body temperature and ambient pressure saturated with water vapor) (see Miller et al., 2005).

[4] The cities/municipalities examined in those seven studies include Guangzhou, Harbin, Shenyang, Shanghai, Xuanwei and Taiwan.

(Jin et al., 1995; Zhou et al., 2000; Jie et al., 2014). For example, Zhou et al. (2014), in a nine-year prospective cohort study among rural residents aged ≥40 in southern China, find a correlation between using clean fuels like biogas (rather than biomass) and better lung function, as well as a decrease in risk for COPD.

2.2. Fuel-based IAP and non-respiratory diseases

A small body of literature for China does examine the association between fuel-based IAP and cardiovascular diseases (CVD), especially high blood pressure. Among these studies, Baumgartner et al. (2011) find that exposure to indoor biomass combustion is associated with an increase in systolic blood pressure (SBP) and diastolic blood pressure (DBP) among rural Chinese women aged >50. They find no such association, however, among younger cohorts aged 25–50. This latter finding is confirmed in Baumgartner et al. (2014) for 280 non-smoking women aged ≥25 living in a rural region of north-western Yunnan province whose exposure to pyrolytic biomass combustion is strongly associated with blood pressure issues.[5] Lee et al. (2012) also find that the use of in-home solid fuel is significantly correlated with an increase in the probabilities of hypertension, coronary heart disease, and diabetes among adults aged ≥18 at Putuo district of Shanghai.[6]

Domestic cooking fuels are also linked to certain poisonous endemics. For instance, in their study of arsenic exposure associated with coal-burning in a small village in Guizhou province, Shraim et al. (2003) find that females have higher dimethylarsinic acid levels but lower percentages of inorganic arsenic and monomethylarsonic acid than males. Several studies also confirm Finkelman et al.'s (1999) finding that domestic coal combustion results in selenium poisoning and potential mercury poisoning (Zheng et al., 2005; Li et al., 2006; Liu et al., 2007a). Li and Ma (2010) also identify a correlation between an increased incidence of CO poisoning and indoor coal burning, inadequate ventilation and/or inappropriate stove use.

In the context of our study, three aspects of the extant literature are particularly relevant: First, the empirical results for China generally suggest that fuel-based IAP has an adverse impact on both respiratory and non-respiratory health outcomes. Yet most existing studies for China cover only certain provinces or cities (see e.g. Peabody et al., 2005), and no studies use nationally representative data. Second, except for Barone-Adesi et al. (2012) and Zhou et al.

[5] Epidemiological studies in Taiwan provide some evidence that pollutant-based IAP might be an emerging risk factor for CVD, including blood pressure and heart rate increments (see Chuang et al., 2010; Chuang et al., 2013; Lin et al., 2009).

[6] It is worth noting that, although Lee et al. (2012) provide evidence for the association between in-home solid fuel use and cardiovascular diseases (CV), measures of CV are self-reported except for blood pressure. Their analysis is also based on cross-sectional data, and reported household solid fuels are used for both cooking and heating.

(2014), who use retrospective data to evaluate the impact of domestic cooking fuels on lung cancer and COPD, respectively, nearly all extant studies use cross-sectional data. Third, only a limited number of studies examine the association between fuel-based IAP and cardiovascular diseases (CVD) in China. Hence, despite recent findings on the association between IAP and blood pressure (see Baumgartner et al., 2011; Baumgartner et al., 2014), relatively little is known about how IAP exposure from domestic cooking fuels is related to other risk predictors of CVD such as inflammation. Furthermore, as emphasized by Baumgartner et al., (2011) and Baumgartner et al., (2014), the evidence on the association between blood pressure (BP) and domestic cooking fuels is exclusively based on cross-sectional data. A major contribution of our study, therefore, is the provision of nationally representative results based on panel data that allow us to control for unobserved heterogeneity. Given that high blood pressure results in around half of all CVD (Mackay and Mensa, 2004)[7], another contribution is our assessment of how domestic cooking fuels affect some major risk predictors of CVD (such as high blood pressure), which may shed light on the mechanisms through which IAP operates on health outcomes.

3. Data and methods

The China Family Panel Studies (CFPS), conducted by Peking University's Institute of Social Science Survey, currently comprises two waves, 2010 and 2012, which cover 25 provinces/municipalities/autonomous regions representing 95% of the Chinese population. Thus, CFPS is a nationally representative dataset. Additionally, the CFPS employs a multistage sampling design and aims to capture socio-economic development, as well as economic and non-economic well-being in Chinese households.[8] The CFPS sample for our study, taken from both waves, includes 12,901 rural women aged ≥16.

The China Health and Nutrition Survey (CHNS), administered in nine waves – 1989, 1991, 1993, 1997, 2000, 2004, 2004, 2009 and 2011 – covers nine provinces (Liaoning, Heilongjiang,[9] Jiangsu, Shandong, Henan, Hubei, Hunan, Guangxi and Guizhou) that are regionally representative but differ in geographic and economic conditions. The multistage random cluster design of this longitudinal dataset allows us to capture the spatio-temporal

[7] It is worth highlighting that every 10-point increase in diastolic blood pressure or every 20 point increase in systolic blood pressure will lead to the doubled risks of cardiovascular diseases (Mackay and Mensa, 2004).

[8] The CFPS excludes Hong Kong, Macao, Taiwan, Xinjiang, Qinghai, Inner Mongolia, Ningxia and Hainan (see Xie et al., 2014, for detailed information about the sampling design).

[9] Heilongjiang province was added as the ninth province in 1997.

dynamics of the social, economic and health situations of the Chinese population.[10] Our CHNS sample, taken from the 1991 to 2009 waves, includes 15,539 rural women aged ≥16.

We use these CFPS and CHNS data to examine the impact of domestic cooking fuels on the health (including self-reported chronic or acute diseases and health status) of rural Chinese women. The advantage of the CFPS is that it is the first dataset that contains nationally representative information on health and cooking fuels. The CHNS, although not nationally representative, has three other advantages that we exploit in the analysis: a long-running panel, information on time exposure to cooking and, particularly important, biomarker information that allows a unique analysis of the association between cooking fuels and certain risk factors of CVDs.

3.1. Health measures

The CFPS captures illness or disease with the following question: *"Have you felt any physical discomfort during the preceding two weeks"*, which is coded 1 if the answer is yes, and 0 otherwise. Similarly, the CHNS captures general health as follows: *"Have you suffered from a chronic or acute disease during the past 4 weeks?*", which is represented by a dummy coded 1 if the answer is yes, and 0 otherwise. The CFPS also assesses self-reported health (SRH) status on a 5-point scale: *"How would you rate your health status? 1 = excellent, 2 = very good, 3 = good, 4 = fair and 5 = poor"*. We rescale these values from 1 = *poor* to 5 = *excellent* to bring them into line with the corresponding 4-point CHNS item, which appears only in the 1997 to 2006 waves: "*Right now, how would you describe your health compared to that of other people your age? 1 = bad; 2 = fair; 3 = good; 4 = excellent*." It should be noted that SRH covers not only mental and physical health but also subjective experience of acute and chronic diseases and overall feelings of well-being (Xie and Mo, 2014).

For the risk predictors of CVD, we use CHNS data on systolic and diastolic blood pressure (SBP/DBP) and inflammation (C-reactive protein, CRP) as proxies. In the CHNS, blood pressure information is available for the 1991 to 2009 waves[11] based on three successive pairs of blood pressure measurements taken by a health professional using a mercury sphygmomanometer at intervals of at least one minute. We calculate the average values of SBP and DBP based on the second and third measurements to overcome potential measurement biases (Lei et al., 2012). We follow the conventional way of dealing with CRP values and

[10] See Zhang et al. (2014) for more detailed information on the CHNS.
[11] See Lei et al. (2012) for a detailed overview of the blood pressure measurements in the CHNS.

remove those with values beyond 10 mg/dl (Pearson et al., 2004). Inflammation is only available in the 2009 CHNS wave and it captured by a dummy variable equal to 1 if the high sensitivity C-reactive protein (CRP) $\geq$ 3 mg/dl and 0 otherwise (Yan et al., 2012) mainly because values above 3 mg/L are associated with high risk of future cardiovascular events (Pearson et al., 2004).

We also test for several other major disease symptoms potentially related to fuel-based IAP using binary dummies. In the CFPS data, these are fever, pain, diarrhoea, cough and palpitation, whereas in the CHNS, they are fever, cough, diarrhoea, asthma, eye disease and heart disease/chest pain.

3.2. Household cooking fuels (HCF)

As in Peabody et al. (2005), we investigate three major household cooking fuels – wood/straw, coal and liquefied natural gas – which account for 84% of total household fuel consumption in the CFPS and 88% in the CHNS. The household cooking fuel variable can thus take three values: 0 = wood/straw, 1 = coal and 2 = liquefied natural gas (LNG), with wood/straw as the reference group. As noted by Peabody et al. (2005), fuel types might be better proxies for pollution exposure than the type of stove used in the household, although both are only indirect measures of exposure to IAP. The fuel-based approach is, however, regarded as the most accurate technique for assessing IAP in developing countries (Mestl and Edwards, 2011).

Regarding possible mechanisms of how different cooking fuels operate on health, exposure to biomass fuel use (like wood/straw) is associated with chronic bronchitis or chronic obstructive pulmonary disease mainly through reduced mucociliary clearance and macrophage response (Bruce et al., 2000). In addition, exposure to biomass smoke for cooking is also a potential risk factor for lung cancer because some carcinogenic compounds are generated via polycyclic aromatic hydrocarbons (Bruce et al., 2000). Use of biomass fuel also leads to cataracts since oxidative changes happen with the absorption of toxins into the lens (Bruce et al., 2000).

As Zhang and Smith (2007) have highlighted, coal contains intrinsic contaminants such as sulphur, and coal burning for cooking generally takes longer than LNG or other gas combustion (Zhang, 1996). More importantly, the concentrations of particulate matter (PM) and sulphur dioxide (SO2) are highest in the kitchen when coal is used for cooking (Zhang and Smith, 2007), thereby giving rise to some cardiovascular diseases (Bruce et al., 2000; Baumgartner et al., 2011).

Lastly, we categorize the independent variables into three subgroups – individual, family and community characteristics – discussed separately below:

3.3. Individual characteristics

Individual controls include age, age squared, educational level, marital status, job status and current smoking behaviour. Education is measured on a 6-point scale recoded as a dummy variable: 1 = illiterate, 2 = primary school, 3 = middle school, 4 = high school, 5 = vocational school and 6 = university or higher. Marital status is a dichotomous variable equal to 1 if the respondent is married and 0 if the respondent is living together with a partner, divorced or widowed. Job status is a dummy variable equal to 1 if the respondent is currently employed and 0 otherwise. Current smoking behaviour is a binary variable equal to 1 if the respondent has smoked in the past month and 0 otherwise. The definitions of the individual variables are the same for both the CFPS and the CHNS except for time spent cooking, which is only available in the CHNS data as measured by the following question: *"During the previous week, how much time (in hours) did you spend per day, on average, cooking food for the household?"*

3.4. Family characteristics

The family controls are household size and household income, which for ease of comparability is adjusted to 2012 in the CFPS and to 2011 in the CHNS. In addition, we also introduce the availability of safe drinking water, existence of flushing toilets, clean trash treatment (for the CFPS), existence of excreta around the living place (for the CHNS) and availability of electricity. Regarding the CFPS, the safe drinking water is defined as 1 if the household's drinking source is tap water or mineral/purified water, 0 otherwise. Flushing toilet facility equals 1 if the household mostly use an indoor/outdoor flushing toilet, 0 otherwise. The electricity variable equals 1 if the household has occasional or no power outage, 0 otherwise. As with the CHNS, safe drinking water variable is a binary variable equal to 1 if the household's water source is a water plant or ground water above 5 meter deep, 0 otherwise. The flushing toilet variable equals 1 if the household can access an in-house/out-house flushing toilet facility, 0 otherwise. Excreta dummy equals 1 if there is no excreta around the dwelling place, 0 otherwise. Availability of electricity is a dummy variable equal to 1 if electric facilities are accessible for the household, 0 otherwise.

3.5. Community characteristics

In the CFPS, we employ the distance in the community to the nearest hospital/medical center, which is a continuous variable measured in kilometres. In the CHNS, the availability of health facilities in the community is defined by a dummy variable equal to 1 if a health facility is located in the village/neighbourhood and 0 if in another village/town/city or in the respondent's city but in a different neighbourhood. Similarly, we also add the distance to the health facility measured in kilometres.

3.6. Estimation approaches

To investigate the association between domestic cooking fuels and general disease (or specific symptoms) in a cross-sectional setting in the CFPS and the CHNS, we adopt a probit regression model of the following form:

$$GD_i = \alpha_0 + \alpha_1 HCF + \alpha_2 I_i + \alpha_3 F + \alpha_4 C + \alpha_5 Y + \alpha_6 P + \varepsilon_i \quad (1)$$

where GD_i is a binary variable indicating general chronic or acute disease or a specific disease symptom of individual *i*, and *HCF* denotes dummies for the three household cooking fuels, wood/straw, coal and liquefied natural gas, with wood/straw as the reference group (applied to both the CFPS and the CHNS). I_i is a vector of individual *i*'s characteristics, *F* is a vector of family characteristics and C is a vector of community characteristics. *Y* is a vector of year dummies (with 2010 and 1991 as the reference year in the CFPS and CHNS, respectively), and *P* is a vector of provincial dummy variables (with Liaoning as the reference province in both datasets). α_1 is the key coefficient of interest, and ε_i is the error term. Because both surveys use a multistage sampling design, we take into account clustering at the village/neighbourhood (CFPS) or community (CHNS) level.

To assess the effects of household cooking fuels on certain risk predictors of cardiovascular diseases (using the CHNS measure of inflammation), we use a specification similar to equation (1) but without year dummies. For SRH (measured on a 4-point scale in the CHNS and a 5-point scale in the CFPS), we estimate the impact of household cooking fuels with an ordered probit model using data from the CFPS and the CHNS and specifications similar to equation (1).

As regards the association between domestic cooking fuels and disease in general (or specific symptoms), since domestic cooking fuels in a household would not change within a short-term period, we account for the potential biases associated with individual unobservables

by estimating the following random effects probit model with data from the CFPS and the CHNS:

$$GD_{it} = I(X'_{it}\beta + F_t + \alpha_i + \varepsilon_{it} \geq 0) \qquad (2)$$

where GD_{it} denotes general chronic or acute disease of individual *i* at time *t*, $I(\cdot)$ is an indicator function that equals 1 if its argument is true (0 otherwise), and X_{it} is a vector of individual characteristics including age and time spent on cooking. F_t captures the two family characteristics, household income and household size, while α_i is assumed to be a random variable.

Because self-reported health is ordinal in both data sets under consideration, for this variable we adopt a random effects ordered probit estimation that uses a specification similar to equation (2):

$$SRH^*_{it} = X_{it}\beta + \alpha_i + \varepsilon_{it} \qquad (3)$$

$$SRH_{it} = \begin{cases} 1, if\ SRH^*_{it} \leq c_1 \\ 2, if\ c_1 < SRH^*_{it} \leq c_2 \\ \vdots \\ K, if\ c_{K-1} < SRH^*_{it} \end{cases} \qquad (4)$$

where SRH^*_{it} (linked to the observed ordinal response categories SRH_{it}) is a latent variable of self-reported health for individual *i* at time *t*. X_{it} represents observed characteristics, α_i is a random variable, *c* is a set of cut-off points $c_1, c_2 \ldots, c_{K-1}$ and *K* is the number of possible outcomes.[12]

We use a balanced panel in the CFPS and an unbalanced panel in the CHNS. In order to test for possible attrition bias in the CHNS, we employ the variable addition test proposed by Verbeek and Nijman (1992). The test adds a variable to the right side of equations (2) and (3) that counts the number of surveyed years that each respondent is present in the survey. The test of attrition bias is evaluated on the significance of this added variable, and insignificance denotes the nonexistence of attrition bias.

To analyse the association between household cooking fuels and SBP or DBP levels in the CHNS data, we employ the following ordinary least squares estimation (OLS):

[12] See Alsakka and Gwilym (2010) for a specific application of the random effects ordered probit approach to examine the sources of heterogeneity in sovereign credit ratings in emerging economies.

$$CVD_i = \beta_0 + \beta_1 HCF + \beta_2 I_i + \beta_3 F + \beta_4 C + \beta_5 P + \beta_6 Y + \varepsilon_i \qquad (5)$$

where CVD_i is a continuous variable denoting the SBP/DBP level of individual *i*, and *HCF* is a dummy for household cooking fuel. I_i is a vector of individual *i*'s characteristics, *F* is a vector of family characteristics and C is a vector of community characteristics. *P* is a vector of provincial dummy variables (with Liaoning as the reference), and *Y* is a vector of year dummies (with 1991 as the reference year). β_1 is the key coefficient of interest and ε_i is the error term. Because the CHNS uses a multistage sampling design, we also take into account clustering at the community level. All the health measures, together with the data and methods used, are summarized in Table 1. In order to take account of individual unobserved heterogeneity, we also estimate a corresponding panel model using data from the CHNS.

Table 1. Summary of health measures, data and methods

Health measures	Description	Definition	Data source	Years	Methodology
Self-reported acute/chronic disease	Have you felt any physical discomfort during the preceding two weeks?	A binary variable equal to 1 if the respondent has felt discomfort, and 0 otherwise.	CFPS	2010-2012	Probit model Random-effect probit model
	Have you suffered from a chronic or acute disease during the past 4 weeks?	A dummy that equals 1 if the respondent has suffered from a chronic or acute disease, and 0 otherwise	CHNS	1991-2009	Probit model Random-effect probit model
Self-reported health (SRH)	How would you rate your health status? 1=excellent, 2=very good, 3=good, 4=fair and 5=poor	A 5-point scale ranging from 1=poor to 5=excellent	CFPS	2010-2012	Ordered probit model Random effect ordered probit model
	Right now, how would you describe your health compared to that of other people your age? 1=bad; 2=fair; 3=good; 4=excellent.	A 4-point scale ranging from 1=bad to 4=excellent	CHNS	1997-2006	Ordered probit model Random effect ordered probit model
Systolic blood pressure (SBP)	Measurements are taken three times by a health professional using a mercury sphygmomanometer	The average value of SBP based on the second and third measurements	CHNS	1991-2009	Ordinary least squares model Fixed effect model
Diastolic blood pressure (DBP)	Measurements are taken three times by a health professional using a mercury sphygmomanometer	The average value of SBP based on the second and third measurements	CHNS	1991-2009	Ordinary least square model Fixed effect model
Inflammation	Using high sensitivity C-reactive protein	A dummy equal to 1 if the high sensitivity C-	CHNS	2009	Probit model

		reactive protein exceeds 3 mg/dl, 0 otherwise			

4. Results

4.1. Descriptive statistics

As Appendix Table A1 shows, the average age of rural women in the CFPS is around 46, which is quite similar to the 45 years in the CHNS (see Table A2). Wood/straw is more predominant as a cooking fuel (CFPS: 63.7%; CHNS: 40.9%) than either coal or LNG (see Appendix Tables A1 and A2). The proportion of LNG, one of the cleaner fuel types, is higher in the CFPS (25.7%) than in the CHNS (17.1%), but this outcome could result from the different periods covered by each survey (2010–2012 and 1991–2009, respectively). The differences between the two datasets could also be attributed to the different geographic regions they cover. Additionally, the trends of domestic cooking fuels from 1991 to 2009 are illustrated in the Appendix Figure A1. Two points are worth highlighting: First, the patterns of domestic cooking fuels have shifted dramatically. Specifically, wood/straw use for household cooking has declined, from 36.9% in 1991 to 17.8% in 2009. Similarly, coal use also has experienced a decrease during the same period, from 36% to 23.2%. However, the use of LNG for domestic cooking has increased significantly, from 15.2% to 22.4%. Second, although the use of wood/straw, coal has declined but LNG has increased steadily during the last decade, those three fuels are still the predominant types for household cooking fuels. Such temporal changes in our sample are very comparable to the results from Duan et al. (2014) based on the data from the CHNS 1991 to 2006.

4.2. Cross-sectional evidence of HCF and health: the CFPS and the CHNS

According to the pooled cross-sectional results for the CFPS (Table 2), women whose households rely on LNG are less likely to suffer from chronic or acute disease than those whose households rely on wood/straw (CFPS: marginal effect (ME) = -4.4%, column 1). Likewise, women in households using either coal or LNG are more likely to self-report better health status than those in households using wood/straw.[13] Nevertheless, the point estimates for LNG are larger than those for coal (2.9% for LNG versus 1.9% for coal, respectively, column 2),

[13] When evaluating the impact of domestic cooking fuels on health outcomes, we also introduce a dummy variable indicating whether any highly polluting enterprises (e.g. a chemical plant) within a 5-kilometer radius centered at the village or residential level exist. The results are quite similar and are available from the authors upon request.

suggesting that cleaner cooking fuels like LNG are beneficial to health. Interestingly, as for the CHNS, we also cannot observe any association between coal/LNG and chronic/acute disease (column 3). Nonetheless, we indeed find that, relative to wood/straw, LNG use is associated with better self-reported health (ME=2.1%, column 4).[14]

Table 2. (Ordered) probit estimates for the effect of household cooking fuels on women's health in rural area: CFPS 2010–2012 and CHNS 1991–2009

Variables	CFPS		CHNS	
	(1) Chronic/acute disease	(2) SRH (excellent)	(3) Chronic/acute disease	(4) SRH (excellent)
Coal	0.010	0.019*	-0.008	0.009
	(0.020)	(0.011)	(0.008)	(0.007)
LNG	-0.044***	0.029***	-0.009	0.021***
	(0.015)	(0.009)	(0.010)	(0.008)
N	12901	12901	15539	8409
Pseudo R^2	0.065	0.147	0.090	0.081

Notes: The dependent variable is a dummy variable for whether the respondent has suffered from a chronic or acute disease (1 = yes, 0 = no) or self-reported health (SRH) measured on a 5-point scale from 1 = poor to 5 = excellent in the CFPS and a 4-point scale from 1 = poor to 4 = excellent in the CHNS. Controls include a dummy for household cooking fuel (0 = wood/straw, 1 = coal, 2 = LNG, with wood/straw as the referent), individual characteristics, family characteristics, community characteristics, provincial dummies (with Liaoning as the referent in both the CFPS and CHNS) and year dummies (with 2010 and 1991 as the referents in the CFPS and CHNS, respectively). Also reported are marginal effects, which for SRH indicate the probability of excellent health. Village/neighbour or community-level clustered standard errors are in parentheses. * p < 0.1, ** p < 0.05, *** p < 0.01.

To check possible effects of domestic cooking fuels on health among different age cohorts, we split the sample into two groups: women aged 16-50 and 50+. The results demonstrate that, relative to wood/straw, women aged 16-50 in households using LNG are more likely to self-report better health in the CFPS. However, this is not the case for those women aged 16-50 in the CHNS. Interestingly, we can consistently find that LNG use for cooking is associated with improved self-rated health among women aged 50+ in both the CFPS and the CHNS (see Table A3 in the Appendix).

4.3. Panel evidence for HCF and health: the CFPS

[14] We also find that, in the CFPS, having a large household and clean rubbish treatment are correlated with a lower probability of chronic/acute disease. Interestingly, having a higher education and household income are related to better health status. However, a longer distance to health facilities is negatively associated with reporting better health. Regarding the CHNS, being employed, having a large household size, better sanitation conditions (e.g. having a flushing toilet in the household) and no excreta around the dwelling place are associated with lower levels of chronic/acute disease. Additionally, having a higher household income and no excreta around the dwelling place are positively related to reporting better health. Also note that age is positively associated with the probability of chronic/acute diseases but negatively correlated with reporting better health in both datasets.

Given the possible existence of individual unobservables or omitted factors, we also estimate random effects (ordered) probit estimates based on the CFPS data, which reveal a significantly negative association between the probability of chronic/acute diseases and cooking with LNG (ME=-5.2%, Table 3, column 1).[15] They also reveal a significantly positive association between self-reported health and cooking with coal or LNG (ME: 2.0% for LNG versus 2.3% for coal, respectively, column 2), which implies that both these fuels are less detrimental than wood.[16]

Table 3. Random effects (ordered) probit estimates for the effect of household cooking fuels on women's health in rural areas: CFPS 2010–2012

Variables	(1) Chronic/acute disease Random effects probit	(2) Self-reported health (excellent) Random effects ordered probit
Coal	-0.006	0.023**
	(0.020)	(0.010)
LNG	-0.052***	0.020***
	(0.015)	(0.007)
N	9770	10000

Notes: The dependent variable is either a dummy for whether the respondent has suffered from a chronic or acute illness (1 = yes, 0 = no) or a 5-point measure of self-reported health problems (from 1 = poor to 5 = excellent). The controls are domestic cooking fuel (0 = wood/straw, 1 = coal, 2 = liquefied natural gas (LNG), with wood/straw as the referent), age, age squared, household income and household size. Also reported are marginal effects, which for SRH indicate the probability of excellent health. Standard errors are in parentheses. * p < 0.1, ** p < 0.05, *** p < 0.01

Our analysis for the two age groups (15-50 and 50+) shows that, compared with wood/straw, LNG use for cooking is strongly associated with better self-reported health among women aged 50+, but not for those aged 16-50. Thus, older rural women are more likely to benefit from positive effects of cleaner fuels like LNG (see Table A4 in the Appendix).

4.4. Panel evidence for HCF and health: the CHNS

We perform a similar panel analysis with the CHNS but introduce an additional control for the time a woman spends cooking (TSC), which is divided into four categories: 1 = TSC<1 hour/day, 2 = 1≤TSC<2 hours/day, 3 = 2≤TSC<3 hours/day, and 4 = TSC≥3 hours/day, with

[15] Note that we also employ a likelihood-ratio test to check the appropriateness of random effects (ordered) probit estimates and results demonstrate that there is enough variability between individuals to favor random effects (ordered) probit estimates instead of (ordered) probit estimates. This is also the case for the CHNS estimation. The results are not reported here, but available from the authors upon request.

[16] It is also interesting to note that, for self-reported health, age is negatively associated with better health while having a higher household income is positively correlated with improved health.

TSC<1 hour/day as the reference.[17] Two of the findings are worth emphasizing: First, the random effects probit estimations (Table 4, column 1) reveal no significantly negative association between domestic LNG use and chronic/acute diseases while the random effects ordered probit estimations (column 2) show a significant positive effect of LNG use on self-reported health (ME=2.2%). Taken together, these results suggest that LNG may lead to better health outcomes. Second, none of the coefficients on time spent cooking are significant (columns 1 and 2), which could imply that the mere use of a specific fuel may lead to IAP persistency. In other words, the spatio-temporal distribution of pollutant concentrations could be affected by house structure, room layout and ventilation (Zhang and Smith, 2007). [18] For the test of attrition bias, the results from random effect probit and order probit estimates indicate that this additional variable is insignificant (not reported here). Thus, it suggests that endogenous attrition bias is not a serious issue in our case.[19]

Table 4. Random effects (ordered) probit estimates for the effect of household cooking fuels on women's health in rural areas: CHNS 1991-2009

Variables	(1) Chronic/acute disease Random effects probit	(2) Self-reported health (excellent) Random effects ordered probit
Coal	-0.007	0.002
	(0.006)	(0.005)
LNG	-0.013	0.022***
	(0.008)	(0.006)
1≤TSC<2 hours/day	-0.003	-0.005
	(0.008)	(0.006)
2≤TSC<3 hours/day	-0.011	0.003
	(0.008)	(0.006)
TSC≥3 hours/day	-0.012	-0.011
	(0.009)	(0.008)
N	10090	7023

Notes: The dependent variable is either a dummy for whether or not the respondent has suffered from a chronic or acute disease or a 4-point measure of self-reported health (1 = poor, 2 = fair, 3 = good and 4 = excellent). The controls are cooking fuel (0 = wood, 1 = coal, 2 = liquefied natural gas (LNG), with wood/straw as the referent), a dummy for time spent cooking (TSC, 1 = TSC<1 hour/day, 2 = 1≤TSC<2 hours/day, 3 = 2≤TSC<3 hours/day, 4 = TSC≥3 hours/day, with group 1 as the referent), age, age squared, household income (inflated to 2011) and household size. Also reported are marginal effects, which for SRH indicate the probability of excellent health. Standard errors are in parentheses. * p < 0.1, ** p < 0.05, *** p < 0.01.

[17] Similar to the results from the CFPS, we also find that age is negatively related to improved health and having a higher level of household income is positively correlated with better health.

[18] We also interact fuel type with TSC, but the estimated coefficients remains insignificant.

[19] To capture some potential effects of seasonal changes for cooking fuel use within a household, we also introduce interviewed month into our estimates. The results, available upon request, are qualitatively very similar to those without controlling for seasonal changes.

To detect the possible heterogeneity of domestic cooking fuel use on various different age groups, we rerun the estimates for rural women aged 15-50 and 50+ separately. The results indicate that in both age groups, relative to wood/straw, LNG use is associated with a higher probability of self-reported improved health. One interesting point to emphasize is that, for those women aged 50+, time spent cooking above 3 hours/day is correlated with a lower probability of reporting better health status, perhaps suggesting that older women are more likely to suffer from longer exposure to IAP (see Table A5 in the Appendix).

4.5. HCF and major risk predictors of cardiovascular diseases (CVD): the CHNS

Even though China is facing a major increase in cardiovascular diseases (Gordon-Larsen et al., 2013; Yan et al., 2012), empirical evidence of the possible effects of household cooking fuels on CVD is quite limited (Fullerton et al., 2008). High blood pressure is one of the major risk contributors for CVD (Mackay and Mensa, 2004). Following Baumgartner et al., (2011), we analyse the levels of systolic blood pressure (SBP) and diastolic blood pressure (DBP). We dichotomize the whole sample into two age groups, namely 16-50 and 50+, because the relative increased risk in cardiovascular diseases begin usually in the fifth decade of life (see for instance Mackay and Mensa, 2004). In our analysis of the CHNS data, we note that, relative to wood/straw, cooking with coal is associated with an increase in raised levels of SBP and DBP among rural women aged 50+ (see columns 2 and 4 in Table 5), but no significance is found for those women aged 16-50. This result supports Baumgartner et al.'s (2011) conclusion that indoor solid fuel combustion is related to a higher level of SBP among women aged 50+ in rural Yunnan, China. We also introduce inflammation as another potential risk factor for CVD. Nevertheless, we find no association between domestic cooking fuels and inflammation (see column 5).

Table 5. OLS/probit estimates for the effect of household cooking fuels on women's blood pressure/inflammation in rural areas: CHNS

	OLS estimates				Probit estimate
Variables	(1) Systolic blood pressure 16-50	(2) Systolic blood pressure 50+	(3) Diastolic blood pressure 16-50	(4) Diastolic blood pressure 50+	(5) Inflammation
Coal	0.070	1.809**	0.520	0.912*	0.016
	(0.589)	(0.747)	(0.390)	(0.464)	(0.032)
LNG	0.335	1.133	0.358	0.484	0.005
	(0.616)	(0.891)	(0.419)	(0.572)	(0.031)

N	6288	9251	6288	9251	1637
Pseudo R^2/ Adjusted R^2	0.108	0.174	0.102	0.093	0.109

Notes: The dependent variable is a dummy for whether the respondent has suffered from inflammation (1 = yes, 0 = no) or from levels of systolic and diastolic blood pressure. The information on blood pressure is available from 1991 to 2009; that on inflammation, only in 2009. The controls include a dummy for household cooking fuel (0 = wood, 1 = coal, 2 = liquefied natural gas, with wood as the referent), individual characteristics, family characteristics, community characteristics, provincial dummies (with Liaoning as the referent) and year dummies (with 1991 as the referent). Marginal effects are reported for the probit estimate of inflammation. Community clustered standard errors are in parentheses. * p < 0.1, ** p < 0.05, *** p < 0.01.

To assess the possible existence of some unobserved or omitted factors, we also estimate a fixed effects model.[20] As Table 6 demonstrates, coal use for domestic cooking is still significantly and positively correlated with an elevated level of systolic blood pressure among women aged 50+ (see column 2), which is quite similar to that of the OLS estimates in the Table 5. Additionally, longer time exposure is also associated with raised levels of SBP among women aged 50+ (>3 hours/day) and DBP among those aged 16-50 (1-3 hours/day) (columns 2 and 3).

Table 6. Fixed effects estimates for the effect of household cooking fuels on women' blood pressure in rural area: CHNS 1991-2009

Variables	Systolic blood pressure		Diastolic blood pressure	
	(1)	(2)	(3)	(4)
	16-50	50+	16-50	50+
Coal	-0.599	2.267*	0.288	0.079
	(0.669)	(1.229)	(0.481)	(0.707)
LNG	-0.131	-0.573	0.654	-0.215
	(0.768)	(1.438)	(0.553)	(0.827)
1-2 hours	0.524	1.536	1.076**	-0.111
	(0.612)	(1.006)	(0.440)	(0.579)
2-3 hours	-0.036	1.731	0.920*	-0.110
	(0.664)	(1.129)	(0.478)	(0.649)
>3 hours	-1.064	2.603**	-0.008	0.417
	(0.726)	(1.300)	(0.523)	(0.748)
N	6252	3838	6252	3838
R^2	0.088	0.102	0.066	0.028

Notes: The dependent variables are levels of systolic/diastolic blood pressure. Controls are cooking fuel choices (0=wood, 1=coal, 2=liquefied natural gas (LNG), wood/straw as the reference), dummies of time spent cooking (1=<1 hour/day, 2=1<= & <2 hours/day, 3=2<= & <3 hours/day, 4=>=3 & <4, and 5=>=4 & <5 hours/day, with group 1 as the reference groups), age, age squared, household income (inflated to 2011) and household size. Standard errors are in parentheses, * p < 0.1, ** p < 0.05, *** p < 0.01.

4.6. HCF and specific symptoms of chronic or acute diseases

[20] We performed a Hausman test to detect the appropriateness of fixed effects over a random effects model. Results are clearly in favor of a fixed effects model.

Finally, as Table 7 shows, we find no significant association between coal/LNG (with wood/straw as the referent) and specific disease symptoms except for significantly negative associations with palpitation in the CFPS, and with eye disease but significantly positive association with heart/chest pain in the CHNS.

Table 7. Probit estimates for the effect of domestic cooking fuels on women's health (specific symptoms) in rural areas: CFPS 2010-2012 & CHNS 1991-2009

Variables	CFPS				
	Fever	Diarrhoea	Cough	Pain	Palpitation
Coal	-0.003	0.003	0.003	0.007	0.006
	(0.005)	(0.002)	(0.004)	(0.015)	(0.005)
LNG	0.001	-0.001	0.003	-0.018	-0.008*
	(0.004)	(0.002)	(0.003)	(0.012)	(0.004)
N	12637	10713	12623	12901	12901
Pseudo R^2	0.023	0.055	0.031	0.059	0.072
Variables	CHNS				
	Fever/cough	Diarrhoea	Asthma	Eye	Heart/chest
Coal	-0.003	0.003	-0.007	-0.003*	0.004*
	(0.005)	(0.003)	(0.007)	(0.002)	(0.002)
LNG	-0.001	-0.002	-0.007	-0.004**	0.003
	(0.006)	(0.004)	(0.010)	(0.002)	(0.002)
N	15539	15539	1078	14299	15539
Pseudo R^2	0.081	0.069	0.297	0.208	0.172

Notes: The dependent variable is a dummy for whether the respondent has suffered from fever, diarrhoea, cough, asthma, pain, palpitation, eye or heart disease/chest pain (1 = yes, 0 = no). Asthma is only available in the year of 2009 for the CHNS. The controls are a dummy for household cooking fuel (0 = wood/straw, 1 = coal, 2 = liquefied natural gas, with wood/straw as the referent), individual characteristics, family characteristics, community characteristics, provincial dummies (with Liaoning as the referent in both the CFPS and CHNS) and year dummies (with 2010 and 1991 as the referent in the CFPS and CHNS, respectively). Marginal effects are reported. Village/neighbour or community-level clustered standard errors are in parentheses, * p < 0.1, ** p < 0.05, *** p < 0.01.

The results from the random effects probit model are illustrated in the Table 8. Two points are worth noting: First, when taking wood/straw as the reference category, we mostly fail to observe any significant association between coal/LNG and specific disease symptoms, which is consistent with the previous cross-sectional results in Table 7. Second, in the CFPS, compared with wood/straw, LNG is associated with a lower probability of diarrhoea (ME=-2.6%). Nonetheless, coal or LNG use for cooking is related to a decreased probability of fever in the CHNS (ME: -1.2% for LNG versus -0.8% for coal, respectively).[21]

[21] When adding the interview month into the estimates, we find that the results are qualitatively similar.

Table 8. Random effects probit estimates for the effect of domestic cooking fuels on women's health (specific symptoms) in rural areas: CFPS 2010-2012 & CHNS 1991-2009

Variables	CFPS				
	Fever	Diarrhoea	Cough	Pain	Palpitation
Coal	-0.002	-0.005	0.003	0.001	0.003
	(0.004)	(0.013)	(0.002)	(0.001)	(0.003)
LNG	0.002	-0.026***	-0.002	0.002	-0.003
	(0.003)	(0.009)	(0.002)	(0.001)	(0.002)
N	10002	10002	10002	10002	10002
			CHNS		
Variables	Fever	Diarrhoea	Eye	Heart/chest pain	
Coal	-0.008*	-0.001	-0.001	0.001	
	(0.004)	(0.002)	(0.001)	(0.001)	
LNG	-0.012**	-0.003	-0.001	0.001	
	(0.006)	(0.003)	(0.001)	(0.001)	
1-2 hours	-0.005	0.003	0.0002	0.0004	
	(0.005)	(0.003)	(0.001)	(0.001)	
2-3 hours	-0.012**	-0.001	-0.001	-0.002	
	(0.006)	(0.003)	(0.001)	(0.002)	
>3 hours	-0.022***	-0.004	-0.0001	-0.0004	
	(0.007)	(0.003)	(0.001)	(0.001)	
N	10090	10090	10090	10090	

Notes: The dependent variable is a dummy for whether the respondent has suffered from fever, diarrhoea, cough, pain, palpitation, eye or heart disease/chest pain (1 = yes, 0 = no). In the CFPS, the controls are a dummy for household cooking fuel (0 = wood/straw, 1 = coal, 2 = liquefied natural gas, with wood/straw as the referent), age, age squared, household income (inflated to 2012) and household size. For the CHNS, controls are similar to those in the CFPS but add dummies of time spent cooking (1=<1 hour/day, 2=1<= & <2 hours/day, 3=2<= & <3 hours/day, 4=>=3 & <4, and 5=>=4 & <5 hours/day. Marginal effects are reported. Standard errors are in parentheses, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

5. Conclusions

This analysis of data from the China Family Panel Studies (CFPS) and the China Health and Nutrition Survey (CHNS) examines the association between the type of domestic cooking fuel and women's health in rural China. Not only does it expand the extant literature by using the most recent nationally and regionally representative datasets within panel settings, it also explores how household cooking fuels are related to the health impairments associated with certain risk predictor for cardiovascular disease. Our results indicate that, relative to those who use traditional biomass fuels like wood/straw, women in rural households who cook with liquefied natural gas are less likely to suffer from chronic or acute diseases and more likely to report better health outcomes. They also associate the use of coal with increased levels of systolic and diastolic blood pressure, suggesting that solid fuel use could be affecting the risk predictors for cardiovascular disease in rural China. This finding is consistent with previous epidemiological studies in rural China (see Baumgartner et al., 2011; Baumgartner et al., 2014).

Such incomplete combustion of solid fuels like biomass and coal emits several pollutants like PM and other toxic compounds (Smith et al., 2004). In particular, PM inhalation results in oxidative stress and systematic inflammation as well (Barregard et al., 2006; Pope and Dockery, 2006), thereby leading to elevated blood pressure levels (Bautista et al., 2005).

Admittedly, however, the observed effects are generally small in magnitude, perhaps because of certain study limitations. One important constraint is that even though a significant proportion of rural households use a mixture of fuels (Zhang and Smith, 2007), our dataset only permits an analysis of primary fuel use, which makes it difficult to isolate the effect of a specific fuel type. Another potential problem is the endogeneity encountered in virtually all studies on this topic (Dulfo et al., 2008), which in the absence of adequate instruments is difficult to solve.[22] Nevertheless, despite these limitations, our analysis does provide evidence for an association between fuel type and health problems among women in rural China. Any such evidence implies that shifting from dirty fuels such as biomass to cleaner choices like liquefied natural gas or electricity could improve health outcomes. This shift, however, is inherently a long-term process, one likely to be hampered by the affordability and availability of cleaner fuels in rural areas (see Baumgartner et al., 2012).

Acknowledgements

For their financial support of the collection and analysis of China Health and Nutrition Survey (CHNS) data files from 1989 to 2011, as well as from future surveys, we thank the National Institute of Nutrition and Food Safety, China Center for Disease Control and Prevention, Carolina Population Center (5 R24 HD050924), the University of North Carolina at Chapel Hill, and the NIH (R01-HD30880, DK056350, R24 HD050924, and R01-HD38700). We further thank the Fogarty International Center, the NIH, and the China-Japan Friendship Hospital, Ministry of Health for their help with the CHNS 2009. We are also very grateful to the Institute of Social Science Survey at Peking University for providing data from the China

[22] We also estimate a special regressor model (SRM) meant for binary choice model estimation when one or more regressors are endogenous or mismeasured (Lewbel et al., 2012). Following Lewbel et al. (2012), we adopt age as the special regressor mainly because it is both continuous and exogenous. In our case, the instrument candidates include 2011-adjusted prices of coal briquettes and liquefied natural gas at the community level. Taking wood/straw and coal as the referent, we do observe a negative yet insignificant association between LNG and chronic/acute diseases. Based on the Hansen J-test and Wald F-test, our instruments appear to be reasonable (results available from the authors upon request). Nevertheless, as Lewbel et al. (2012) emphasize, such an instrumental technique can be problematic if the association between special regressors and other endogenous variables leads to violation of the special regressor's conditional independence from the error term. In addition, the fact that the model imposes fewer assumptions on error term distribution leads to larger standard errors (Lewbel et al., 2012).

Family Panel Studies (CFPS). This analysis is an output of a scholarship from the Food Security Center at the University of Hohenheim, which is part of the DAAD (German Academic Exchange Service) “Exceed” programme and supported by DAAD and the German Federal Ministry for Economic Cooperation and Development (BMZ). This paper was presented at the 2014 Global Food Studies Seminar in Adelaide, the 2015 XVIII Applied Economics Meeting in Alicante, the 2015 ESRC Multi-disciplinary Health & Biomarkers Conference in Essex, and the 11th World Congress on Health Economics in Milan. We would like to thank the participants, as well as Jan Michael Bauer, Meena Kumari and Henk Folmer for valuables comments on an earlier version of this paper. The usual disclaimer applies.

References

Alsakka, R. and Gwilym, O. (2010), 'A random effects ordered probit model for rating migrations', *Finance Research Letters* **7**(3): 140-147.

Balakrishnan, K., Sankar, S., Ghosh, S., Thangavel, G., Mukhopadhyay, K., Ramaswamy, P., Johnson, P. and Thanasekaraan, V. (2014), 'Household Air Pollution Related to Solid Cookfuel Use: The Exposure and Health Situation in Developing Countries', In: D. Barceló & A.G. Kostianoy (eds.), *The Handbook of Environmental Chemistry*, Heidelberg: Springer. pp. 1–20.

Barone-Adesi, F., Chapman, R. S., Silverman, D. T., He, X., Hu, W., Vermeulen, R., Ning, B., Fraumeni, J. F., Jr., Rothman, N. and Lan, Q. (2012), 'Risk of lung cancer associated with domestic use of coal in Xuanwei, China: retrospective cohort study', *BMJ* **345**: e5414.

Baumgartner, J., Schauer, J. J., Ezzati, M., Lu, L., Cheng, C., Patz, J. A. and Bautista, L. E. (2011), 'Indoor air pollution and blood pressure in adult women living in rural China', *Environmental Health Perspectives* **119**(10): 1390-1395.

Baumgartner, J., Smith, K. R. and Chockalingam, A. (2012), 'Reducing CVD Through Improvements in Household Energy', *Global Heart* **7**(3): 243-247.

Baumgartner, J., Zhang, Y., Schauer, J. J., Huang, W., Wang, Y. and Ezzati, M. (2014), 'Highway proximity and black carbon from cookstoves as a risk factor for higher blood pressure in rural China', *Proceedings of the National Academy of Sciences* **111**(36): 13229-13234.

Bentayeb, M., Simoni, M., Norback, D., Baldacci, S., Maio, S., Viegi, G. and Annesi-Maesano, I. (2013), 'Indoor air pollution and respiratory health in the elderly', *Journal of Environmental Science and Health. Part A, Toxic/Hazardous Substances & Environmental Engineering* **48**(14): 1783-1789.

Bruce, N., Perez-Padilla, R. and Albalak, R. (2000), 'Indoor air pollution in developing countries: a major environmental and public health challenge', *Bulletin of the World Health Organization*, **78**(9): 1078-1092.

Barregard, L., Sällsten, G., Gustafson, P., Andersson, L., Johansson, L., Basu, S. and Stigendal, L. (2006), 'Experimental exposure to wood smoke particles in healthy humans: effects on markers of inflammation, coagulation and lipid peroxidation', *Inhalation Toxicology* **18**(11): 845-853.

Bautista, L.E., Vera, L.M., Arenas, I.A., Gamarra, G. (2005), 'Independent association between inflammatory markers (C-reactive protein, interleukin-6, and TNF-α) and essential hypertension', *Journal of Human Hypertension* **19**:149-154.

Chen, W., Zheng, R., Zeng, H. and Zhang, S. (2014), 'The epidemiology of lung cancer', *Journal of Cancer Biology & Research* **2**(1): 1043.

Chuang, K. J., Yan, Y. H. and Cheng, T. J. (2010), 'Effect of air pollution on blood pressure, blood lipids, and blood sugar: A population-based approach', *Journal of Occupational and Environmental Medicine* **52**(3): 258-262.

Chuang, W. Y., Lin, L. Y. and Chuang, K. J. (2013), 'Indoor air quality and cardiovascular effects in the Taiwanese general population: Generalized additive mixed models analysis', Paper presented at the World Congress on Engineering & Computer Science 2013, 3-5 July 2013, London, U.K.

Duan, X., Jiang, Y., Wang, B., Zhao, X., Shen, G., Cao, S., Huang, N., Qian, Y., Chen, Y. and Wang, L. (2014), 'Household fuel use for cooking and heating in China: Results from the first Chinese Environmental Exposure-Related Human Activity Patterns Survey (CEERHAPS)', *Applied Energy* **136**: 692-703.

Dulfo, E., Greenstone, M. and Hanna, R. (2008), 'Indoor air pollution, health and economic well-being', *Surveys and Perspectives Integrating Environment & Society* **1**: 1-9.

Finkelman, R. B., Belkin, H. E. and Zheng, B. (1999), 'Health impacts of domestic coal use in China', *Proceedings of the National Academy of Sciences* **96**(7): 3427-3431.

Fullerton, D. G., Bruce, N. and Gordon, S. B. (2008), 'Indoor air pollution from biomass fuel smoke is a major health concern in the developing world', *Transactions of the Royal Society of Tropical Medicine and Hygiene* **102**(9): 843-851.

Jie, Y., Houjin, H., Xun, M., Kebin, L., Xuesong, Y. and Jie, X. (2014), 'Relationship between pulmonary function and indoor air pollution from coal combustion among adult residents in an inner-city area of southwest China', *Brazilian Journal of Medical and Biological Research* **47**(11): 982-989.

Jin, Y., Zhou, X. and He, X. (1995), 'Study on effect of indoor pollutants of lung functionalities from combustion of fuel', *China Public Health* **11**: 18-19.

Jin, Z. Y., Wu, M., Han, R. Q., Zhang, X. F., Wang, X. S., Liu, A. M., Zhou, J. Y., Lu, Q. Y., Kim, C. H., Mu, L., Zhang, Z. F. and Zhao, J. K. (2014), 'Household ventilation may reduce effects of indoor air pollutants for prevention of lung cancer: A case-control study in a Chinese population', *PLoS One* **9**(7): e102685.

Kan, X., Chiang, C. Y., Enarson, D. A., Chen, W., Yang, J. and Chen, G. (2011), 'Indoor solid fuel use and tuberculosis in China: A matched case-control study', *BMC Public Health* **11**: 498.

Kim, C., Gao, Y. T., Xiang, Y. B., Barone-Adesi, F., Zhang, Y., Hosgood, H. D., Ma, S., Shu, X. O., Ji, B. T., Chow, W. H., Seow, W. J., Bassig, B., Cai, Q., Zheng, W., Rothman, N. and Lan, Q. (2014), 'Home kitchen ventilation, cooking fuels, and lung cancer risk in a prospective cohort of never smoking women in Shanghai, China', *International Journal of Cancer* **136**(3): 632-638.

Kurmi, O. P., Arya, P. H., Lam, K. B., Sorahan, T. and Ayres, J. G. (2012), 'Lung cancer risk and solid fuel smoke exposure: A systematic review and meta-analysis', *The European Respiratory Journal* **40**(5): 1228-1237.

Lei, X., Yin, N. and Zhao, Y. (2012), 'Socioeconomic status and chronic diseases: The case of hypertension in China', *China Economic Review* **23**(1): 105-121.

Lewbel, A., Dong, Y. Y. and Yang, T. (2012), 'Comparing features of convenient estimators for binary choice models with endogenous regressors', *Canadian Journal of Economics* **45**(3): 809-829.

Li, C. and Ma, H. L. (2010), 'Risk factors of non-occupational carbon monoxide poisoning during the 2008 ice storm in Guiyang county, Hunan province, China', *Public Health Reports* **125**(4): 605-610.

Li, D. S., An, D., Zhou, Y. S., Liu, J. and Waalkes, M. P. (2006), 'Current status and prevention strategy for coal-arsenic poisoning in Guizhou, China', *Journal of Health, Population and Nutrition* **24**(3): 273-276.

Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H., AlMazroa, M. A., Amann, M., Anderson, H. R., Andrews, K. G., et al. (2012), 'A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions,

1990–2010: A systematic analysis for the Global Burden of Disease Study 2010', *The Lancet* **380**(9859): 2224-2260.

Lin, H. H., Suk, C. W., Lo, H. L., Huang, R. Y., Enarson, D. A. and Chiang, C. Y. (2014), 'Indoor air pollution from solid fuel and tuberculosis: A systematic review and meta-analysis', *The International Journal of Tuberculosis and Lung Disease* **18**(5): 613-621.

Lee, M. S., Hang, J. Q., Zhang, F. Y., Dai, H. L., Su, L. and Christiani, D. C. (2012), 'In-home solid fuel use and cardiovascular disease: a cross-sectional analysis of the Shanghai Putuo study', *Environmental Health* **11**:18.

Lin, L. Y., Lin, C. Y., Lin, Y. C. and Chuang, K. J. (2009), 'The effects of indoor particles on blood pressure and heart rate among young adults in Taipei, Taiwan', *Indoor Air* **19**(6): 482-488.

Liu, G., Zheng, L., Duzgoren-Aydin, N., Gao, L., Liu, J. and Peng, Z. (2007a), 'Health Effects of Arsenic, Fluorine, and Selenium from Indoor Burning of Chinese Coal', in Whitacre, D., Ware, G., Nigg, H., Doerge, D., Albert, L., de Voogt, P., Gerba, C., Hutzinger, O., Knaak, J., Mayer, F., Morgan, D. P., Park, D., Tjeerdema, R., Yang, R. H. and Gunther, F. (eds.), *Reviews of Environmental Contamination and Toxicology*, New York: Springer, pp. 89-106.

Liu, S., Zhou, Y., Wang, X., Wang, D., Lu, J., Zheng, J., Zhong, N. and Ran, P. (2007b), 'Biomass fuels are the probable risk factor for chronic obstructive pulmonary disease in rural South China', *Thorax* **62**(10): 889-897.

Mestl, H. E. and Edwards, R. (2011), 'Global burden of disease as a result of indoor air pollution in Shaanxi, Hubei and Zhejiang, China', *Science of the Total Environment* **409**(8): 1391-1398.

Miller, M. R., Hankinson, J., Brusasco, V., Burgos, F., Casaburi, R., Coates, A., Crapo, R., Enright, P., van der Grinten, C. P., Gustafsson, P., et al. (2005), 'Standardisation of spirometry', *The European Respiratory Journal* **26**(2): 319-338.

Mackay, J., Mensa, G.A. (2004), 'The Atlas of Heart Disease and Stroke', Geneva: World Health Organization.

Pan, X. C., Dong, Z., Wang, L. and Yue, W. (2002), 'An evaluation of the indoor/outdoor air pollution and respiratory health of farmers living in rural areas of Anhui province, China', in Proceedings, Indoor Air 2002, Monterey: International Academy of Indoor Air Sciences.

Peabody, J. W., Riddell, T. J., Smith, K. R., Liu, Y., Zhao, Y., Gong, J., Milet, M. and Sinton, J. E. (2005), 'Indoor air pollution in rural China: Cooking fuels, stoves, and health status', *Archives of Environment & Occupational Health* **60**(2): 86-95.

Pearson, T.A., Mensah, G.A., Hong, Y. and Smith, S.C. (2004), CDC/AHA workshop on markers of inflammation and cardiovascular disease: Application to clinical and public health practice: overview', *Circulation* **110**(25): e543-e544.

Pope, C.A. III. and Dockery, D.W. (2006), 'Health effects of fine particulate air pollution: Lines that connect', *Journal of the Air & Waste Management Association* **56**(6): 709-742.

Reid, B. C., Ghazarian, A. A., DeMarini, D. M., Sapkota, A., Jack, D., Lan, Q., Winn, D. M. and Birnbaum, L. S. (2012), 'Research opportunities for cancer associated with indoor air pollution from solid-fuel combustion', *Environmental Health Perspectives* **120**(11): 1495-1498.

She, J., Yang, P., Hong, Q. and Bai, C. (2013), 'Lung cancer in China: Challenges and interventions', *Chest* **143**(4): 1117-1126.

Shraim, A., Cui, X., Li, S., Ng, J. C., Wang, J., Jin, Y., Liu, Y., Guo, L., Li, D., Wang, S., Zhang, R. and Hirano, S. (2003), 'Arsenic speciation in the urine and hair of individuals exposed to airborne arsenic through coal-burning in Guizhou, PR China', *Toxicology Letters* **137**(1–2): 35-48.

Smith, K. R., Mehta, S. and Maeusezahl-Feuz (2004), 'Indoor air pollution from household use of solid fuels', in Ezzati, M., Lopez, A. D., Rodgers, A. and Murray, C. J. L. (eds.), *Comparative Quantification of Health Risks*, Geneva: World Health Organization, pp. 1435-1493.

United Nations Development Group (2013), *A Million Voices: The World We Want*, New York: United Nations Development Group.

Verbeek, M. and Nijman, T. (1992), 'Testing for Selectivity Bias in Panel Data Models', *International Economic Review* **33**(3): 681-703.

World Health Organization and United Nations Development Programme (2009), '*The Energy Access Situation in Developing Countries*', New York: United Nations Development Programme.

Xie, S. and Mo, T. (2014), 'The impact of education on health in China', *China Economic Review* **29**: 1-18.

Xie, Y., Hu, J. and Zhang, C. (2014), 'The China Family Panel Studies (CFPS): Design and practice', *Chinese Journal of Sociology (Shehui)* **34**: 1-33.

Xu, F., Yin, X., Shen, H., Xu, Y., Ware, R. S. and Owen, N. (2007), 'Better understanding the influence of cigarette smoking and indoor air pollution on chronic obstructive pulmonary disease: A case-control study in Mainland China', *Respirology* **12**(6): 891-897.

Yan, S., Li, J., Li, S., Zhang, B., Du, S., Gordon-Larsen, P., Adair, L. and Popkin, B. (2012), 'The expanding burden of cardiometabolic risk in China: The China Health and Nutrition Survey', *Obesity Reviews* **13**(9): 810-821.

Zhang, B., Zhai, F. Y., Du, S. F. and Popkin, B. M. (2014), 'The China Health and Nutrition Survey, 1989-2011', *Obesity Reviews* **15 Suppl 1**: 2-7.

Zhang, J. J. and Smith, K. R. (2007), 'Household air pollution from coal and biomass fuels in China: Measurements, health impacts, and interventions', *Environmental Health Perspectives* **115**(6): 848-855.

Zhang, Q., Li, L., Smith, M., Guo, Y., Whitlock, G., Bian, Z., Kurmi, O., Collins, R., Chen, J., Lv, S., Pang, Z., Chen, C., Chen, N., Xiong, Y., Peto, R., Chen, Z. and China Kadoorie Biobank study, c. (2013), 'Exhaled carbon monoxide and its associations with smoking, indoor household air pollution and chronic respiratory diseases among 512,000 Chinese adults', *International Journal of Epidemiology* **42**(5): 1464-1475.

Zhang, Y. B. and Wu, Y. (2012), *China - Health Impacts of Indoor Air Pollution : At A Glance*, East Asia and Pacific (EAP) Clean Stove Initiative Knowledge Exchange Series, Washington, DC: World Bank.

Zhao, Y., Wang, S., Aunan, K., Seip, H. M. and Hao, J. (2006), 'Air pollution and lung cancer risks in China-a meta-analysis', *Science of the Total Environment* **366**(2-3): 500-513.

Zheng, B. S., Wang, B. B., Ding, Z. H., Zhou, D. X., Zhou, Y. S., Chen, Z., Chen, C. C. and Finkelman, R. B. (2005), 'Endemic arsenosis caused by indoor combustion of high-as coal in Guizhou Province, P.R. China', *Environmental Geochemistry and Health* **27**(5-6): 521-528.

Zhou, B. S., Wang, T. J., Guan, P. and Wu, J. M. (2000), 'Indoor air pollution and pulmonary adenocarcinoma among females: A case-control study in Shenyang, China', *Oncology Reports* **7**(6): 1253-1262.

Zhou, Y., Zou, Y., Li, X., Chen, S., Zhao, Z., He, F., Zou, W., Luo, Q., Li, W., Pan, Y., Deng, X., Wang, X., Qiu, R., Liu, S., Zheng, J., Zhong, N. and Ran, P. (2014), 'Lung function and incidence of chronic obstructive pulmonary disease after improved cooking fuels and kitchen ventilation: A 9-year prospective cohort study', *PLoS Medicine* **11**(3): e1001621.

Zhang, J., Wang, L., Liu, J., Zhou, Y., Wen, T. and Meng, D. (1996), '24-hour NO_2 level in kitchen air and 24-hour NO_2 personal exposure level and its effect on urinary HOP', *Environmental Health* **13**:193–196. (in Chinese)

Appendix:

Table A1. Descriptive statistics: CFPS 2010–2012

Variables	Obs	Mean	Std. Dev.	Min	Max
Dependent variables					
Chronic/acute disease	12901	0.347	0.476	0	1
Fever	12901	0.022	0.147	0	1
Pain	12901	0.156	0.363	0	1
Diarrhoea	12901	0.005	0.072	0	1
Cough	12901	0.015	0.121	0	1
Palpitation	12901	0.033	0.177	0	1
Self-reported health (SRH)					
Poor	12901	0.146	0.353	0	1
Fair	12901	0.154	0.361	0	1
Good	12901	0.171	0.376	0	1
Very good	12901	0.262	0.440	0	1
Excellent	12901	0.267	0.443	0	1
Household cooking fuels					
Wood/straw	12901	0.637	0.481	0	1
Coal	12901	0.106	0.308	0	1
Liquefied natural gas (LNG)	12901	0.257	0.437	0	1
Individual characteristics					
Age	12901	46.059	16.053	16	97
Working status	12901	0.490	0.500	0	1
Education levels					
Illiterate	12901	0.516	0.500	0	1
Primary school	12901	0.220	0.414	0	1
Middle school	12901	0.194	0.395	0	1
High school	12901	0.052	0.223	0	1
Vocational school	12901	0.012	0.110	0	1
University or higher	12901	0.005	0.073	0	1
Marital status	12901	0.834	0.372	0	1
Currently smoking	12901	0.039	0.195	0	1
Family characteristics					
Household income (log)	12901	9.877	1.148	0.693	14.253
Household size	12901	4.674	1.922	1	26
Drinking water	12901	0.421	0.494	0	1
Electricity	12901	0.944	0.230	0	1
Flushing toilet	12901	0.209	0.407	0	1
Clean trash treatment	12901	0.148	0.355	0	1
Community Characteristics					
Distance to the health facility (km)	12901	1.247	1.670	0.001	9.500

Notes: Self-reported health (SRH) is measured on a 5-point-scale (1 = poor, 2 = fair, 3= good, 4 = very good and 5 = excellent). The household cooking fuels are wood/straw, coal and liquefied natural gas (0 = wood/straw, 1 = coal, 2 = liquefied natural gas (LNG)). The education level dummy is measured on a 6-point scale (1= illiterate, 2 = primary school, 3 = middle school, 4 = high school, 5 = vocational school and 6 = university or higher). Dummies are also included for marital status (1 = married, 0 = others), working status (1 = currently employed, 0 = currently unemployed) and smoking behaviour at the time of interview (1 if the respondent had smoked for the past month, 0 otherwise).
Source: China Family Panel Studies 2010 and 2012.

Table A2. Descriptive statistics: CHNS 1991-2009

Variables	Obs	Mean	Std. Dev.	Min	Max
Dependent variables					
Chronic/acute disease	15539	0.109	0.311	0	1
Fever	15539	0.046	0.210	0	1
Diarrhoea	15539	0.017	0.130	0	1
Asthma	1078	0.001	0.031	0	1
Eye disease	15539	0.004	0.066	0	1
Heart disease/chest pain	15539	0.009	0.092	0	1
Self-reported health (SRH)					
Bad	8409	0.070	0.256	0	1
Fair	8409	0.317	0.465	0	1
Good	8409	0.492	0.500	0	1
Excellent	8409	0.121	0.326	0	1
Household cooking fuels					
Wood/straw	15539	0.409	0.492	0	1
Coal	15539	0.420	0.494	0	1
Liquefied natural gas (LNG)	15539	0.171	0.377	0	1
Individual characteristics					
Age	15539	44.959	15.728	16	97.84
Working status	15539	0.705	0.456	0	1
Education levels					
Illiterate	15539	0.300	0.458	0	1
Primary school	15539	0.337	0.473	0	1
Middle school	15539	0.263	0.440	0	1
High school	15539	0.073	0.260	0	1
Vocational school	15539	0.020	0.140	0	1
University or higher	15539	0.007	0.083	0	1
Marital status	15539	0.799	0.400	0	1
Currently smoking	15539	0.037	0.189	0	1
Time spent cooking (hours/day)	10227	1.728	0.932	0.017	4.667
Family characteristics					
Household income (log)	15539	9.390	1.011	1.156	13.414
Household size	15539	4.198	1.591	1	13
Water	15539	0.827	0.378	0	1
Flushing toilet	15539	0.207	0.405	0	1
No excreta around the dwelling place	15539	0.568	0.495	0	1
Electricity	15539	0.984	0.127	0	1
Community characteristics					
Location of health facility	15539	0.752	0.432	0	1
Distance to the health facility (km)	15539	0.923	3.573	0	60

Notes: Self-reported health (SRH), which is only available from 1997 to 2006, is measured on a 4-point scale (1 = poor, 2 = fair, 3 = good and 4 = excellent). Asthma is only available in the year of 2009. The household cooking fuels are wood, coal and liquefied natural gas (0 = wood, 1 = coal, 2 = liquefied natural gas (LNG)). The education level dummy is measured on a 6-point scale (0 = illiterate, 1 = primary school, 2 = middle school, 3 = high school, 4 = vocational school and 5 = university or higher). Dummies are also included for marital status (1 = married, 0 = other), working status (1 = currently employed, 0 = currently unemployed) and smoking behaviour at the time of interview (1 if the respondent was currently smoking cigarettes, 0 otherwise).
Source: China Health and Nutrition Survey 1991, 1993, 1997, 2000, 2004, 2006 and 2009.

Table A3. Ordered probit estimates for cooking fuel choice on women self-reported health in rural area: CFPS 2010-2012 and CHNS 1991-2009

Variables	CFPS		CHNS	
	(1)	(2)	(3)	(4)
	16-50	50+	16-50	50+
Coal	0.010	0.027**	0.016	0.003
	(0.015)	(0.012)	(0.012)	(0.006)
LNG	0.025**	0.035***	0.014	0.022***
	(0.012)	(0.011)	(0.013)	(0.007)
N	7958	4945	5107	3302
Pseudo R^2	0.140	0.109	0.052	0.042

Notes: The dependent variable is self-reported health (SRH) measured on a 5-point scale from 1 = poor to 5 = excellent in the CFPS or a 4-point scale from 1 = poor to 4 = excellent in the CHNS. Controls include dummies of household cooking fuel choices (0=wood, 1=coal, 2=liquefied natural gas, wood as the reference), individual characteristics (including age, age squared, education level, marital status, job status, current smoking behavior, participation into food preparation and cooking), family characteristics (translog household income inflated to 2011, household size, the availability of safe drinking water, sanitation and clean trash treatment), community characteristics (distance to the health facility), provincial dummies (Liaoning as the reference) and year dummies (2010 as the reference). Also reported are marginal effects, which for SRH indicate the probability of excellent health. Village/neighbour clustered standard errors are in parentheses, * p < 0.1, ** p < 0.05, *** p < 0.01.

Table A4. Random effects ordered probit estimates for cooking fuel choice on women self-reported health in rural area: CFPS 2010-2012

Variables	Random effects ordered probit:	Random effects ordered probit:
	16-50	50+
Coal	0.021	0.024*
	(0.016)	(0.013)
LNG	0.012	0.027***
	(0.011)	(0.009)
N	6011	3989

Notes: The dependent variable is self-reported health (SRH) measured on a 5-point-scale variable of self-reported health status (from 1=very unhealthy to 5=very healthy). Controls include cooking fuel choices (0=wood, 1=coal, 2=liquefied natural gas, wood as the reference), age, age squared, household income (inflated to 2011) and household size. Also reported are marginal effects, which for SRH indicate the probability of excellent health. Standard errors are in parentheses, * p < 0.1, ** p < 0.05, *** p < 0.01.

Table A5. Random effects ordered probit estimates for household cooking fuels on women self-reported health in rural area: CHNS 1991-2009

Variables	Random effects ordered probit: 16-50	Random effects ordered probit: 50+
Coal	-0.001	0.005
	(0.009)	(0.004)
LNG	0.020*	0.020***
	(0.011)	(0.006)
1-2 hours	0.001	-0.007
	(0.010)	(0.005)
2-3 hours	0.007	-0.001
	(0.011)	(0.005)
>3 hours	-0.001	-0.014**
	(0.013)	(0.007)
N	4085	2938

Notes: The dependent variable is self-reported health (SRH) measured on a 4-point scale from 1 = poor to 4 = excellent. Controls include cooking fuel choices (0=wood, 1=coal, 2=liquefied natural gas (LNG), wood/straw as the reference), dummies of time spent cooking (1=<1 hour/day, 2=1<= & <2 hours/day, 3=2<= & <3 hours/day, 4=>=3 & <4, and 5=>=4 & <5 hours/day, with group 1 as the reference groups), age, age squared, household income (inflated to 2011) and household size. Also reported are marginal effects, which for SRH indicate the probability of excellent health. Standard errors are in parentheses, * p < 0.1, ** p < 0.05, *** p < 0.01.



Figure A1. Temporal change of domestic cooking fuels from 1991 to 2009, China Health and Nutrition Survey

Notes: LNG denotes liquefied natural gas.

Hohenheim Discussion Papers in Business, Economics and Social Sciences

The Faculty of Business, Economics and Social Sciences continues since 2015 the established “FZID Discussion Paper Series” of the “Centre for Research on Innovation and Services (FZID)” under the name “Hohenheim Discussion Papers in Business, Economics and Social Sciences”.

Institutes

510 Institute of Financial Management
520 Institute of Economics
530 Institute of Health Care & Public Management
540 Institute of Communication Science
550 Institute of Law and Social Sciences
560 Institute of Economic and Business Education
570 Institute of Marketing & Management
580 Institute of Interorganisational Management & Performance

Download Hohenheim Discussion Papers in Business, Economics and Social Sciences
from our homepage: https://wiso.uni-hohenheim.de/papers

Nr.	Autor	Titel	Inst.
01-2015	Thomas Beissinger, Philipp Baudy	THE IMPACT OF TEMPORARY AGENCY WORK ON TRADE UNION WAGE SETTING: A Theoretical Analysis	520
02-2015	Fabian Wahl	PARTICIPATIVE POLITICAL INSTITUTIONS AND CITY DEVELOPMENT 800-1800	520
03-2015	Tommaso Proietti, Martyna Marczak, Gianluigi Mazzi	$\text{EURO}\text{MI}_{\text{IND}}$-D: A DENSITY ESTIMATE OF MONTHLY GROSS DOMESTIC PRODUCT FOR THE EURO AREA	520
04-2015	Thomas Beissinger, Nathalie Chusseau, Joël Hellier	OFFSHORING AND LABOUR MARKET REFORMS: MODELLING THE GERMAN EXPERIENCE	520
05-2015	Matthias Mueller, Kristina Bogner, Tobias Buchmann, Muhamed Kudic	SIMULATING KNOWLEDGE DIFFUSION IN FOUR STRUCTURALLY DISTINCT NETWORKS – AN AGENT-BASED SIMULATION MODEL	520
06-2015	Martyna Marczak, Thomas Beissinger	BIDIRECTIONAL RELATIONSHIP BETWEEN INVESTOR SENTIMENT AND EXCESS RETURNS: NEW EVIDENCE FROM THE WAVELET PERSPECTIVE	520
07-2015	Peng Nie, Galit Nimrod, Alfonso Sousa-Poza	INTERNET USE AND SUBJECTIVE WELL-BEING IN CHINA	530
08-2015	Fabian Wahl	THE LONG SHADOW OF HISTORY ROMAN LEGACY AND ECONOMIC DEVELOPMENT – EVIDENCE FROM THE GERMAN LIMES	520
09-2015	Peng Nie, Alfonso Sousa-Poza	COMMUTE TIME AND SUBJECTIVE WELL-BEING IN URBAN CHINA	530

Nr.	Autor	Titel	Inst.
10-2015	Kristina Bogner	THE EFFECT OF PROJECT FUNDING ON INNOVATIVE PERFORMANCE AN AGENT-BASED SIMULATION MODEL	520
11-2015	Bogang Jun, Tai-Yoo Kim	A NEO-SCHUMPETERIAN PERSPECTIVE ON THE ANALYTICAL MACROECONOMIC FRAMEWORK: THE EXPANDED REPRODUCTION SYSTEM	520
12-2015	Volker Grossmann Aderonke Osikominu Marius Osterfeld	ARE SOCIOCULTURAL FACTORS IMPORTANT FOR STUDYING A SCIENCE UNIVERSITY MAJOR?	520
13-2015	Martyna Marczak Tommaso Proietti Stefano Grassi	A DATA–CLEANING AUGMENTED KALMAN FILTER FOR ROBUST ESTIMATION OF STATE SPACE MODELS	520
14-2015	Carolina Castagnetti Luisa Rosti Marina Töpfer	THE REVERSAL OF THE GENDER PAY GAP AMONG PUBLIC-CONTEST SELECTED YOUNG EMPLOYEES	520
15-2015	Alexander Opitz	DEMOCRATIC PROSPECTS IN IMPERIAL RUSSIA: THE REVOLUTION OF 1905 AND THE POLITICAL STOCK MARKET	520
01-2016	Michael Ahlheim, Jan Neidhardt	NON-TRADING BEHAVIOUR IN CHOICE EXPERIMENTS	520
02-2016	Bogang Jun, Alexander Gerybadze, Tai-Yoo Kim	THE LEGACY OF FRIEDRICH LIST: THE EXPANSIVE REPRODUCTION SYSTEM AND THE KOREAN HISTORY OF INDUSTRIALIZATION	520
03-2016	Peng Nie, Alfonso Sousa-Poza	FOOD INSECURITY AMONG OLDER EUROPEANS: EVIDENCE FROM THE SURVEY OF HEALTH, AGEING, AND RETIREMENT IN EUROPE	530
04-2016	Peter Spahn	POPULATION GROWTH, SAVING, INTEREST RATES AND STAGNATION. DISCUSSING THE EGGERTSSON-MEHROTRA-MODEL	520
05-2016	Vincent Dekker, Kristina Strohmaier, Nicole Bosch	A DATA-DRIVEN PROCEDURE TO DETERMINE THE BUNCHING WINDOW – AN APPLICATION TO THE NETHERLANDS	520
06-2016	Philipp Baudy, Dario Cords	DEREGULATION OF TEMPORARY AGENCY EMPLOYMENT IN A UNIONIZED ECONOMY: DOES THIS REALLY LEAD TO A SUBSTITUTION OF REGULAR EMPLOYMENT?	520
07-2016	Robin Jessen, Davud Rostam-Afschar, Sebastian Schmitz	HOW IMPORTANT IS PRECAUTIONARY LABOR SUPPLY?	520
08-2016	Peng Nie, Alfonso Sousa-Poza, Jianhong Xue	FUEL FOR LIFE: DOMESTIC COOKING FUELS AND WOMEN'S HEALTH IN RURAL CHINA	530

FZID Discussion Papers

(published 2009-2014)

Competence Centers

IK	Innovation and Knowledge
ICT	Information Systems and Communication Systems
CRFM	Corporate Finance and Risk Management
HCM	Health Care Management
CM	Communication Management
MM	Marketing Management
ECO	Economics

Download FZID Discussion Papers from our homepage: https://wiso.uni-hohenheim.de/archiv_fzid_papers

Nr.	Autor	Titel	CC
01-2009	Julian P. Christ	NEW ECONOMIC GEOGRAPHY RELOADED: Localized Knowledge Spillovers and the Geography of Innovation	IK
02-2009	André P. Slowak	MARKET FIELD STRUCTURE & DYNAMICS IN INDUSTRIAL AUTOMATION	IK
03-2009	Pier Paolo Saviotti, Andreas Pyka	GENERALIZED BARRIERS TO ENTRY AND ECONOMIC DEVELOPMENT	IK
04-2009	Uwe Focht, Andreas Richter and Jörg Schiller	INTERMEDIATION AND MATCHING IN INSURANCE MARKETS	HCM
05-2009	Julian P. Christ, André P. Slowak	WHY BLU-RAY VS. HD-DVD IS NOT VHS VS. BETAMAX: THE CO-EVOLUTION OF STANDARD-SETTING CONSORTIA	IK
06-2009	Gabriel Felbermayr, Mario Larch and Wolfgang Lechthaler	UNEMPLOYMENT IN AN INTERDEPENDENT WORLD	ECO
07-2009	Steffen Otterbach	MISMATCHES BETWEEN ACTUAL AND PREFERRED WORK TIME: Empirical Evidence of Hours Constraints in 21 Countries	HCM
08-2009	Sven Wydra	PRODUCTION AND EMPLOYMENT IMPACTS OF NEW TECHNOLOGIES – ANALYSIS FOR BIOTECHNOLOGY	IK
09-2009	Ralf Richter, Jochen Streb	CATCHING-UP AND FALLING BEHIND KNOWLEDGE SPILLOVER FROM AMERICAN TO GERMAN MACHINE TOOL MAKERS	IK

Nr.	Autor	Titel	CC
10-2010	Rahel Aichele, Gabriel Felbermayr	KYOTO AND THE CARBON CONTENT OF TRADE	ECO
11-2010	David E. Bloom, Alfonso Sousa-Poza	ECONOMIC CONSEQUENCES OF LOW FERTILITY IN EUROPE	HCM
12-2010	Michael Ahlheim, Oliver Frör	DRINKING AND PROTECTING – A MARKET APPROACH TO THE PRESERVATION OF CORK OAK LANDSCAPES	ECO
13-2010	Michael Ahlheim, Oliver Frör, Antonia Heinke, Nguyen Minh Duc, and Pham Van Dinh	LABOUR AS A UTILITY MEASURE IN CONTINGENT VALUATION STUDIES – HOW GOOD IS IT REALLY?	ECO
14-2010	Julian P. Christ	THE GEOGRAPHY AND CO-LOCATION OF EUROPEAN TECHNOLOGY-SPECIFIC CO-INVENTORSHIP NETWORKS	IK
15-2010	Harald Degner	WINDOWS OF TECHNOLOGICAL OPPORTUNITY DO TECHNOLOGICAL BOOMS INFLUENCE THE RELATIONSHIP BETWEEN FIRM SIZE AND INNOVATIVENESS?	IK
16-2010	Tobias A. Jopp	THE WELFARE STATE EVOLVES: GERMAN KNAPPSCHAFTEN, 1854-1923	HCM
17-2010	Stefan Kirn (Ed.)	PROCESS OF CHANGE IN ORGANISATIONS THROUGH eHEALTH	ICT
18-2010	Jörg Schiller	ÖKONOMISCHE ASPEKTE DER ENTLOHNUNG UND REGULIERUNG UNABHÄNGIGER VERSICHERUNGSVERMITTLER	HCM
19-2010	Frauke Lammers, Jörg Schiller	CONTRACT DESIGN AND INSURANCE FRAUD: AN EXPERIMENTAL INVESTIGATION	HCM
20-2010	Martyna Marczak, Thomas Beissinger	REAL WAGES AND THE BUSINESS CYCLE IN GERMANY	ECO
21-2010	Harald Degner, Jochen Streb	FOREIGN PATENTING IN GERMANY, 1877-1932	IK
22-2010	Heiko Stüber, Thomas Beissinger	DOES DOWNWARD NOMINAL WAGE RIGIDITY DAMPEN WAGE INCREASES?	ECO
23-2010	Mark Spoerer, Jochen Streb	GUNS AND BUTTER – BUT NO MARGARINE: THE IMPACT OF NAZI ECONOMIC POLICIES ON GERMAN FOOD CONSUMPTION, 1933-38	ECO

Nr.	Autor	Titel	CC
24-2011	Dhammika Dharmapala, Nadine Riedel	EARNINGS SHOCKS AND TAX-MOTIVATED INCOME-SHIFTING: EVIDENCE FROM EUROPEAN MULTINATIONALS	ECO
25-2011	Michael Schuele, Stefan Kirn	QUALITATIVES, RÄUMLICHES SCHLIEßEN ZUR KOLLISIONSERKENNUNG UND KOLLISIONSVERMEIDUNG AUTONOMER BDI-AGENTEN	ICT
26-2011	Marcus Müller, Guillaume Stern, Ansger Jacob and Stefan Kirn	VERHALTENSMODELLE FÜR SOFTWAREAGENTEN IM PUBLIC GOODS GAME	ICT
27-2011	Monnet Benoit, Patrick Gbakoua and Alfonso Sousa-Poza	ENGEL CURVES, SPATIAL VARIATION IN PRICES AND DEMAND FOR COMMODITIES IN CÔTE D'IVOIRE	ECO
28-2011	Nadine Riedel, Hannah Schildberg-Hörisch	ASYMMETRIC OBLIGATIONS	ECO
29-2011	Nicole Waidlein	CAUSES OF PERSISTENT PRODUCTIVITY DIFFERENCES IN THE WEST GERMAN STATES IN THE PERIOD FROM 1950 TO 1990	IK
30-2011	Dominik Hartmann, Atilio Arata	MEASURING SOCIAL CAPITAL AND INNOVATION IN POOR AGRICULTURAL COMMUNITIES. THE CASE OF CHÁPARRA - PERU	IK
31-2011	Peter Spahn	DIE WÄHRUNGSKRISENUNION DIE EURO-VERSCHULDUNG DER NATIONALSTAATEN ALS SCHWACHSTELLE DER EWU	ECO
32-2011	Fabian Wahl	DIE ENTWICKLUNG DES LEBENSSTANDARDS IM DRITTEN REICH – EINE GLÜCKSÖKONOMISCHE PERSPEKTIVE	ECO
33-2011	Giorgio Triulzi, Ramon Scholz and Andreas Pyka	R&D AND KNOWLEDGE DYNAMICS IN UNIVERSITY-INDUSTRY RELATIONSHIPS IN BIOTECH AND PHARMACEUTICALS: AN AGENT-BASED MODEL	IK
34-2011	Claus D. Müller-Hengstenberg, Stefan Kirn	ANWENDUNG DES ÖFFENTLICHEN VERGABERECHTS AUF MODERNE IT SOFTWAREENTWICKLUNGSVERFAHREN	ICT
35-2011	Andreas Pyka	AVOIDING EVOLUTIONARY INEFFICIENCIES IN INNOVATION NETWORKS	IK
36-2011	David Bell, Steffen Otterbach and Alfonso Sousa-Poza	WORK HOURS CONSTRAINTS AND HEALTH	HCM
37-2011	Lukas Scheffknecht, Felix Geiger	A BEHAVIORAL MACROECONOMIC MODEL WITH ENDOGENOUS BOOM-BUST CYCLES AND LEVERAGE DYNAMICS	ECO
38-2011	Yin Krogmann, Ulrich Schwalbe	INTER-FIRM R&D NETWORKS IN THE GLOBAL PHARMACEUTICAL BIOTECHNOLOGY INDUSTRY DURING 1985–1998: A CONCEPTUAL AND EMPIRICAL ANALYSIS	IK

Nr.	Autor	Titel	CC
39-2011	Michael Ahlheim, Tobias Börger and Oliver Frör	RESPONDENT INCENTIVES IN CONTINGENT VALUATION: THE ROLE OF RECIPROCITY	ECO
40-2011	Tobias Börger	A DIRECT TEST OF SOCIALLY DESIRABLE RESPONDING IN CONTINGENT VALUATION INTERVIEWS	ECO
41-2011	Ralf Rukwid, Julian P. Christ	QUANTITATIVE CLUSTERIDENTIFIKATION AUF EBENE DER DEUTSCHEN STADT- UND LANDKREISE (1999-2008)	IK

Nr.	Autor	Titel	CC
42-2012	Benjamin Schön, Andreas Pyka	A TAXONOMY OF INNOVATION NETWORKS	IK
43-2012	Dirk Foremny, Nadine Riedel	BUSINESS TAXES AND THE ELECTORAL CYCLE	ECO
44-2012	Gisela Di Meglio, Andreas Pyka and Luis Rubalcaba	VARIETIES OF SERVICE ECONOMIES IN EUROPE	IK
45-2012	Ralf Rukwid, Julian P. Christ	INNOVATIONSPOTENTIALE IN BADEN-WÜRTTEMBERG: PRODUKTIONSCLUSTER IM BEREICH „METALL, ELEKTRO, IKT“ UND REGIONALE VERFÜGBARKEIT AKADEMISCHER FACHKRÄFTE IN DEN MINT-FÄCHERN	IK
46-2012	Julian P. Christ, Ralf Rukwid	INNOVATIONSPOTENTIALE IN BADEN-WÜRTTEMBERG: BRANCHENSPEZIFISCHE FORSCHUNGS- UND ENTWICKLUNGSAKTIVITÄT, REGIONALES PATENTAUFKOMMEN UND BESCHÄFTIGUNGSSTRUKTUR	IK
47-2012	Oliver Sauter	ASSESSING UNCERTAINTY IN EUROPE AND THE US - IS THERE A COMMON FACTOR?	ECO
48-2012	Dominik Hartmann	SEN MEETS SCHUMPETER. INTRODUCING STRUCTURAL AND DYNAMIC ELEMENTS INTO THE HUMAN CAPABILITY APPROACH	IK
49-2012	Harold Paredes-Frigolett, Andreas Pyka	DISTAL EMBEDDING AS A TECHNOLOGY INNOVATION NETWORK FORMATION STRATEGY	IK
50-2012	Martyna Marczak, Víctor Gómez	CYCLICALITY OF REAL WAGES IN THE USA AND GERMANY: NEW INSIGHTS FROM WAVELET ANALYSIS	ECO
51-2012	André P. Slowak	DIE DURCHSETZUNG VON SCHNITTSTELLEN IN DER STANDARDSETZUNG: FALLBEISPIEL LADESYSTEM ELEKTROMOBILITÄT	IK
52-2012	Fabian Wahl	WHY IT MATTERS WHAT PEOPLE THINK - BELIEFS, LEGAL ORIGINS AND THE DEEP ROOTS OF TRUST	ECO
53-2012	Dominik Hartmann, Micha Kaiser	STATISTISCHER ÜBERBLICK DER TÜRKISCHEN MIGRATION IN BADEN-WÜRTTEMBERG UND DEUTSCHLAND	IK
54-2012	Dominik Hartmann, Andreas Pyka, Seda Aydin, Lena Klauß, Fabian Stahl, Ali Santircioglu, Silvia Oberegelsbacher, Sheida Rashidi, Gaye Onan and Suna Erginkoç	IDENTIFIZIERUNG UND ANALYSE DEUTSCH-TÜRKISCHER INNOVATIONSNETZWERKE. ERSTE ERGEBNISSE DES TGIN-PROJEKTES	IK
55-2012	Michael Ahlheim, Tobias Börger and Oliver Frör	THE ECOLOGICAL PRICE OF GETTING RICH IN A GREEN DESERT: A CONTINGENT VALUATION STUDY IN RURAL SOUTHWEST CHINA	ECO

Nr.	Autor	Titel	CC
56-2012	Matthias Strifler Thomas Beissinger	FAIRNESS CONSIDERATIONS IN LABOR UNION WAGE SETTING – A THEORETICAL ANALYSIS	ECO
57-2012	Peter Spahn	INTEGRATION DURCH WÄHRUNGSUNION? DER FALL DER EURO-ZONE	ECO
58-2012	Sibylle H. Lehmann	TAKING FIRMS TO THE STOCK MARKET: IPOS AND THE IMPORTANCE OF LARGE BANKS IN IMPERIAL GERMANY 1896-1913	ECO
59-2012	Sibylle H. Lehmann, Philipp Hauber and Alexander Opitz	POLITICAL RIGHTS, TAXATION, AND FIRM VALUATION – EVIDENCE FROM SAXONY AROUND 1900	ECO
60-2012	Martyna Marczak, Víctor Gómez	SPECTRAN, A SET OF MATLAB PROGRAMS FOR SPECTRAL ANALYSIS	ECO
61-2012	Theresa Lohse, Nadine Riedel	THE IMPACT OF TRANSFER PRICING REGULATIONS ON PROFIT SHIFTING WITHIN EUROPEAN MULTINATIONALS	ECO

Nr.	Autor	Titel	CC
62-2013	Heiko Stüber	REAL WAGE CYCLICALITY OF NEWLY HIRED WORKERS	ECO
63-2013	David E. Bloom, Alfonso Sousa-Poza	AGEING AND PRODUCTIVITY	HCM
64-2013	Martyna Marczak, Víctor Gómez	MONTHLY US BUSINESS CYCLE INDICATORS: A NEW MULTIVARIATE APPROACH BASED ON A BAND-PASS FILTER	ECO
65-2013	Dominik Hartmann, Andreas Pyka	INNOVATION, ECONOMIC DIVERSIFICATION AND HUMAN DEVELOPMENT	IK
66-2013	Christof Ernst, Katharina Richter and Nadine Riedel	CORPORATE TAXATION AND THE QUALITY OF RESEARCH AND DEVELOPMENT	ECO
67-2013	Michael Ahlheim, Oliver Frör, Jiang Tong, Luo Jing and Sonna Pelz	NONUSE VALUES OF CLIMATE POLICY - AN EMPIRICAL STUDY IN XINJIANG AND BEIJING	ECO
68-2013	Michael Ahlheim, Friedrich Schneider	CONSIDERING HOUSEHOLD SIZE IN CONTINGENT VALUATION STUDIES	ECO
69-2013	Fabio Bertoni, Tereza Tykvová	WHICH FORM OF VENTURE CAPITAL IS MOST SUPPORTIVE OF INNOVATION? EVIDENCE FROM EUROPEAN BIOTECHNOLOGY COMPANIES	CFRM
70-2013	Tobias Buchmann, Andreas Pyka	THE EVOLUTION OF INNOVATION NETWORKS: THE CASE OF A GERMAN AUTOMOTIVE NETWORK	IK
71-2013	B. Vermeulen, A. Pyka, J. A. La Poutré and A. G. de Kok	CAPABILITY-BASED GOVERNANCE PATTERNS OVER THE PRODUCT LIFE-CYCLE	IK
72-2013	Beatriz Fabiola López Ulloa, Valerie Møller and Alfonso Sousa-Poza	HOW DOES SUBJECTIVE WELL-BEING EVOLVE WITH AGE? A LITERATURE REVIEW	HCM
73-2013	Wencke Gwozdz, Alfonso Sousa-Poza, Lucia A. Reisch, Wolfgang Ahrens, Stefaan De Henauw, Gabriele Eiben, Juan M. Fernández-Alvira, Charalampos Hadjigeorgiou, Eva Kovács, Fabio Lauria, Toomas Veidebaum, Garrath Williams, Karin Bammann	MATERNAL EMPLOYMENT AND CHILDHOOD OBESITY – A EUROPEAN PERSPECTIVE	HCM

Nr.	Autor	Titel	CC
74-2013	Andreas Haas, Annette Hofmann	RISIKEN AUS CLOUD-COMPUTING-SERVICES: FRAGEN DES RISIKOMANAGEMENTS UND ASPEKTE DER VERSICHERBARKEIT	HCM
75-2013	Yin Krogmann, Nadine Riedel and Ulrich Schwalbe	INTER-FIRM R&D NETWORKS IN PHARMACEUTICAL BIOTECHNOLOGY: WHAT DETERMINES FIRM'S CENTRALITY-BASED PARTNERING CAPABILITY?	ECO, IK
76-2013	Peter Spahn	MACROECONOMIC STABILISATION AND BANK LENDING: A SIMPLE WORKHORSE MODEL	ECO
77-2013	Sheida Rashidi, Andreas Pyka	MIGRATION AND INNOVATION – A SURVEY	IK
78-2013	Benjamin Schön, Andreas Pyka	THE SUCCESS FACTORS OF TECHNOLOGY-SOURCING THROUGH MERGERS & ACQUISITIONS – AN INTUITIVE META-ANALYSIS	IK
79-2013	Irene Prostolupow, Andreas Pyka and Barbara Heller-Schuh	TURKISH-GERMAN INNOVATION NETWORKS IN THE EUROPEAN RESEARCH LANDSCAPE	IK
80-2013	Eva Schlenker, Kai D. Schmid	CAPITAL INCOME SHARES AND INCOME INEQUALITY IN THE EUROPEAN UNION	ECO
81-2013	Michael Ahlheim, Tobias Börger and Oliver Frör	THE INFLUENCE OF ETHNICITY AND CULTURE ON THE VALUATION OF ENVIRONMENTAL IMPROVEMENTS – RESULTS FROM A CVM STUDY IN SOUTHWEST CHINA –	ECO
82-2013	Fabian Wahl	DOES MEDIEVAL TRADE STILL MATTER? HISTORICAL TRADE CENTERS, AGGLOMERATION AND CONTEMPORARY ECONOMIC DEVELOPMENT	ECO
83-2013	Peter Spahn	SUBPRIME AND EURO CRISIS: SHOULD WE BLAME THE ECONOMISTS?	ECO
84-2013	Daniel Guffarth, Michael J. Barber	THE EUROPEAN AEROSPACE R&D COLLABORATION NETWORK	IK
85-2013	Athanasios Saitis	KARTELLBEKÄMPFUNG UND INTERNE KARTELLSTRUKTUREN: EIN NETZWERKTHEORETISCHER ANSATZ	IK

Nr.	Autor	Titel	CC
86-2014	Stefan Kirn, Claus D. Müller-Hengstenberg	INTELLIGENTE (SOFTWARE-)AGENTEN: EINE NEUE HERAUSFORDERUNG FÜR DIE GESELLSCHAFT UND UNSER RECHTSSYSTEM?	ICT
87-2014	Peng Nie, Alfonso Sousa-Poza	MATERNAL EMPLOYMENT AND CHILDHOOD OBESITY IN CHINA: EVIDENCE FROM THE CHINA HEALTH AND NUTRITION SURVEY	HCM
88-2014	Steffen Otterbach, Alfonso Sousa-Poza	JOB INSECURITY, EMPLOYABILITY, AND HEALTH: AN ANALYSIS FOR GERMANY ACROSS GENERATIONS	HCM
89-2014	Carsten Burhop, Sibylle H. Lehmann-Hasemeyer	THE GEOGRAPHY OF STOCK EXCHANGES IN IMPERIAL GERMANY	ECO
90-2014	Martyna Marczak, Tommaso Proietti	OUTLIER DETECTION IN STRUCTURAL TIME SERIES MODELS: THE INDICATOR SATURATION APPROACH	ECO
91-2014	Sophie Urmetzer, Andreas Pyka	VARIETIES OF KNOWLEDGE-BASED BIOECONOMIES	IK
92-2014	Bogang Jun, Joongho Lee	THE TRADEOFF BETWEEN FERTILITY AND EDUCATION: EVIDENCE FROM THE KOREAN DEVELOPMENT PATH	IK
93-2014	Bogang Jun, Tai-Yoo Kim	NON-FINANCIAL HURDLES FOR HUMAN CAPITAL ACCUMULATION: LANDOWNERSHIP IN KOREA UNDER JAPANESE RULE	IK
94-2014	Michael Ahlheim, Oliver Frör, Gerhard Langenberger and Sonna Pelz	CHINESE URBANITES AND THE PRESERVATION OF RARE SPECIES IN REMOTE PARTS OF THE COUNTRY – THE EXAMPLE OF EAGLEWOOD	ECO
95-2014	Harold Paredes-Frigolett, Andreas Pyka, Javier Pereira and Luiz Flávio Autran Monteiro Gomes	RANKING THE PERFORMANCE OF NATIONAL INNOVATION SYSTEMS IN THE IBERIAN PENINSULA AND LATIN AMERICA FROM A NEO-SCHUMPETERIAN ECONOMICS PERSPECTIVE	IK
96-2014	Daniel Guffarth, Michael J. Barber	NETWORK EVOLUTION, SUCCESS, AND REGIONAL DEVELOPMENT IN THE EUROPEAN AEROSPACE INDUSTRY	IK

IMPRINT

University of Hohenheim
Dean’s Office of the Faculty of Business, Economics and Social Sciences
Palace Hohenheim 1 B
70593 Stuttgart | Germany
Fon +49 (0)711 459 22488
Fax +49 (0)711 459 22785
E-mail wiso@uni-hohenheim.de
Web www.wiso.uni-hohenheim.de